

A PRIORI ESTIMATES FOR THE SCALAR CURVATURE EQUATION ON S^3

MATTHIAS SCHNEIDER

ABSTRACT. We obtain a priori estimates for solutions to the prescribed scalar curvature equation on S^3 . The usual non-degeneracy assumption on the curvature function is replaced by a new condition, which is necessary and sufficient for the existence of a priori estimates, when the curvature function is a positive Morse function.

1. INTRODUCTION

Let $N \geq 3$ and S^N be the standard sphere with round metric g_0 induced by $S^N = \partial B_1(0) \subset \mathbb{R}^{N+1}$. We study the problem: Which functions K on S^N occur as scalar curvature of metrics g conformally equivalent to g_0 ? Writing $g = \varphi^{4/(N-2)}g_0$ this is equivalent to solving (see [3])

$$-\frac{4(N-1)}{N-2}\Delta_{S^N}\varphi + N(N-1)\varphi = K(\theta)(\varphi)^{\frac{N+2}{N-2}}, \quad \varphi > 0 \quad \text{in } S^N. \quad (1.1)$$

In stereographic coordinates $\mathcal{S}_\theta(\cdot)$ centered at some point $\theta \in S^N$ equation (1.1) is equivalent to

$$-\Delta u = \frac{K \circ \mathcal{S}_\theta(x)}{N(N-1)} u^{\frac{N+2}{N-2}}, \quad u > 0 \quad \text{in } \mathbb{R}^N, \quad (1.2)$$

where

$$u(x) = \mathcal{R}_\theta(\varphi)(x) := (N(N-2))^{\frac{N-2}{4}} (1 + |x|^2)^{-\frac{N-2}{2}} \varphi \circ \mathcal{S}_\theta(x). \quad (1.3)$$

Obviously, to solve (1.1) the function K has to be positive somewhere. Moreover, there are the *Kazdan-Warner* obstructions [7, 16], if φ solves (1.1) then

$$\int_{S^N} \nabla x_j \cdot \nabla K \varphi^{\frac{2N}{N-2}} = 0 \quad \text{for } j = 1 \dots N+1.$$

In particular, a monotone function of x_1 can not be realized as the scalar curvature of a metric conformal to g_0 .

Numerous studies have been made on equation (1.1) and various sufficient conditions for its solvability have been found (see [2, 4, 6, 11, 12, 18, 19] and the reference therein), usually under a non-degeneracy assumption on K . On S^3 a positive function K is non-degenerate, if

$$\Delta_{S^3} K(\theta) \neq 0 \quad \text{if } \nabla K(\theta) = 0. \quad (\text{nd})$$

Date: December 1, 2004.

1991 *Mathematics Subject Classification.* 35J60, 35J20, 55C21.

Key words and phrases. prescribed scalar curvature, a priori estimates.

For positive Morse functions K on S^3 it is shown in [5, 10, 21] that (1.1) is solvable if K satisfies (nd) and

$$d := -\left(1 + \sum_{\substack{\nabla K(\theta)=0, \\ \Delta_{S^3}K(\theta)<0}} (-1)^{\text{ind}(\theta)}\right) \neq 0, \quad (1.4)$$

where $\text{ind}(\theta)$ is the Morse index of K at θ . We are interested in the case when $N = 3$ and the non-degeneracy assumption (nd) is not satisfied.

To obtain the existence result Bahri and Coron [5] use a detailed analysis of the gradient flow of (1.1) and Schoen and Zhang [21] approximate (1.1) by subcritical problems $p \nearrow \frac{N+2}{N-2}$, which are always solvable, and analyze the possible blow-up of solutions.

We follow the approach suggested in [10] and use a continuity method. We join the curvature function K to the constant function $K_0 \equiv 6$ by a one parameter family $K_t(\theta) := 6(1 + tk(\theta))$, where $k(\theta) := \frac{1}{6}(K(\theta) - 6)$, and consider

$$-8\Delta_{S^3}\varphi + 6\varphi = 6(1 + tk(\theta))\varphi^5, \quad \varphi > 0 \text{ in } S^3, \quad (1.5)$$

or in stereographic coordinates using (1.3) and $k_\theta(x) := k \circ \mathcal{S}_\theta(x)$

$$-\Delta u = (1 + tk_\theta(x))u^5 \text{ in } \mathbb{R}^3, \quad u > 0. \quad (1.6)$$

In general there are no a priori L^∞ -estimates for (1.5) or (1.1) due to the noncompact group of conformal transformations of S^N acting on solutions: the solutions of (1.2) for $k \equiv 0$ form a noncompact manifold (see [9, 14])

$$Z := \left\{ z_{\mu,y}(x) := \mu^{-\frac{N-2}{2}} (N(N-2))^{\frac{N-2}{4}} \left(1 + \left| \frac{x-y}{\mu} \right|^2 \right)^{-\frac{N-2}{2}} \right. \\ \left. : y \in \mathbb{R}^N, \mu > 0 \right\},$$

where $z_{\mu,y}(y) \rightarrow \infty$ as $\mu \rightarrow 0$.

Chang, Gursky and Yang [10] show that if $K \in C^2(S^3)$ is positive and satisfies (nd) then for every $\delta > 0$ there is a constant $C = C(\delta, K) > 0$ such that for all $t \in [\delta, 1]$ and solutions φ_t of (1.5) we have

$$C^{-1} \leq \varphi_t(\theta) \leq C \text{ and } \|\varphi_t\|_{C^{2,\alpha}(S^3)} \leq C.$$

Furthermore, they compute the Leray-Schauder degree for (1.5) for $t > 0$ small, and show that it equals d in (1.4) if K is a Morse function. The a priori estimate implies the invariance of the degree as the parameter t moves to 1 and gives a solution to (1.5) if $d \neq 0$. Chen and Lin [11] show that if $K \in C^2(S^3)$ is a non-degenerate Morse function then C may be chosen independently of $\delta > 0$.

Hence, if (nd) fails, we face two problems: Is the a priori bound still valid and how do critical points of K with $\Delta_{S^3}K = 0$ occur in the index count condition (1.4). Here, we will mainly deal with the question about the a priori bound of solutions.

In the following, unless otherwise stated, we will always assume $N = 3$ and that the function $K \in C^5(S^3)$ is positive. To give our main results we need the following notation. For $k \in C^5(S^3)$ we write $k_\theta = k \circ \mathcal{S}_\theta$ and for a critical

point θ of k we let

$$\begin{aligned} a_0(\theta) &:= \oint_{\mathbb{R}^3} \left(k_\theta(x) - \sum_{\ell=0}^2 \frac{1}{\ell!} D^\ell k_\theta(0)(x)^\ell \right) |x|^{-6}, \\ a_1(\theta) &:= \Delta^2 k_\theta(0) + \nabla(\Delta k_\theta(0)) \cdot (D^2 k_\theta(0))^{-1} \nabla(\Delta k_\theta(0)), \\ a_2(\theta) &:= k_\theta(0) a_1(\theta) - \frac{15}{8\pi} \int_{\partial B_1(0)} |D^2 k_\theta(0)(x)^2|^2, \end{aligned} \quad (1.7)$$

where all differentiations are done in $\mathbb{R}^3$ and $\oint$ is the Cauchy principal value of the integral,

$$\oint_{\mathbb{R}^3} f(x) := \lim_{r \rightarrow 0} \int_{\mathbb{R}^3 \setminus B_r(0)} f(x).$$

Theorem 1.1. *Suppose $1 + k \in C^5(S^3)$ is positive and satisfies*

$D^2 k_\theta(0)$ is invertible, if $\theta \in \mathcal{A} := \{\theta \in S^3 : \nabla k(\theta) = 0 \text{ and } \Delta_{S^3} k(\theta) = 0\}$.

Thus, $\mathcal{A}$ is discrete and hence finite. Denote by M the finite set

$$M := \{\theta \in S^3 : \theta \in \mathcal{A}, a_0(\theta) = 0, \text{ and } a_2(\theta) \neq 0\}.$$

Then for every $\delta > 0$ there is a constant $C = C(k, \delta) > 0$ such that for all

$$t \in (0, 1] \setminus \bigcup_{\theta \in M} B_\delta \left(-\frac{a_1(\theta)}{a_2(\theta)} \right)$$

and solutions φ_t of (1.5) we have

$$C^{-1} \leq \varphi_t(x) \leq C \quad \text{and} \quad \|\varphi_t(x)\|_{C^{2,\alpha}(S^3)} \leq C.$$

Theorem 1.1 extends the known a priori estimates to the case when (nd) may fail. If k satisfies (nd) the solutions are uniformly bounded with respect to $t \in (0, 1]$. Moreover, we get uniform estimates for $t \in (0, 1]$ if $M^* = \emptyset$, where

$$M^* := \{\theta \in M : 0 \leq -a_1(\theta)/a_2(\theta) \leq 1\}.$$

Our results are optimal since we construct for every $\theta \in M^*$ solutions φ_t which blow up as $t \rightarrow -a_1(\theta)/a_2(\theta)$. We say that (t_i, φ_i) *blow up* at the *blow-up point* $\theta \in S^3$, if φ_i solves (1.5) with $t = t_i$, the sequence (t_i) is bounded, and there is (θ_i) converging to θ such that $\varphi_i(\theta_i) \rightarrow \infty$.

Theorem 1.2. *Under the assumptions of Theorem 1.1 let*

$$M_+^* := \{\theta \in M : 0 < -a_1(\theta)/a_2(\theta) \leq 1\}.$$

Then there is $\delta > 0$ such that for any $\theta \in M_+^$ there exists a unique C^1 -curve*

$$\{0 < \mu < \delta\} \ni \mu \mapsto (t^\theta(\mu), \varphi^\theta(\mu, \cdot)) \in (\delta, 1 + \delta) \times C^{2,\alpha}(S^3),$$

such that as $\mu \rightarrow 0$

$$t^\theta(\mu) = -\frac{a_1(\theta)}{a_2(\theta)} + O(\mu^{\frac{1}{4}}),$$

and $\varphi^\theta(\mu, \cdot)$ solves (1.5) for $t = t^\theta(\mu)$ and blows up like

$$\|\mathcal{R}_\theta(\varphi^\theta(\mu, x)) - (1 + t^\theta(\mu)k(\theta))^{-\frac{1}{4}} z_{\mu,0}(x)\|_{\mathcal{D}^{1,2}(\mathbb{R}^3) \cap C^2(B_1(0))} = O(\mu^2).$$

The curves are unique, in the sense that, if $(t_i, \varphi_i) \in (\delta, 1 + \delta) \times C^{2,\alpha}(S^3)$ blow up at some $\theta \in S^3$ then $\theta \in M_+^*$ and there is a sequence of positive numbers (μ_i) converging to zero such that $(t_i, \varphi_i) = (t^\theta(\mu_i), \varphi^\theta(\mu_i, \cdot))$ for all but finitely many $i \in \mathbb{N}$.

Hence, for Morse functions we obtain

Corollary 1.3. *Suppose $1 + k \in C^5(S^3)$ is a positive Morse function. There exists $\delta_0 > 0$, such that for any $0 < \delta < \delta_0$ the solutions of (1.5) are uniformly bounded for $t \in [\delta, 1 + \delta]$, if and only if $M_+^* = \emptyset$.*

For $\theta \in M$ with $a_1(\theta) = 0$ there is always the trivial curve of solutions,

$$\mu \mapsto (0, (\mathcal{R}_\theta)^{-1} z_{\mu,0}) \in \mathbb{R} \times C^{2,\alpha}(S^3),$$

which blow up at θ as $\mu \rightarrow 0$. In order to find a nontrivial curve, i.e. $t(\mu) \in \mathbb{R} \setminus \{0\}$, we need to consider

$$\begin{aligned} a_3(\theta) := & \frac{12}{\pi^2} (D^2 k_\theta(0))^{-1} \nabla(\Delta k_\theta(0)) \cdot \oint_{\mathbb{R}^3} \left(\nabla k_\theta(x) - T_{\nabla k_\theta,0}^2(x) \right) |x|^{-6} \\ & + \frac{48}{\pi^2} (D^2 k_\theta(0))^{-1} \nabla(\Delta k_\theta(0)) \cdot \oint_{\mathbb{R}^3} \left(k_\theta(x) - T_{k_\theta,0}^3(x) \right) \frac{x_i}{|x|^8} \\ & - \frac{120}{\pi^2} \oint_{\mathbb{R}^3} \left(k_\theta(x) - T_{k_\theta,0}^4(x) \right) \frac{1}{|x|^8}, \end{aligned} \quad (1.8)$$

where we abbreviate the m th Taylor polynomial of k in y by

$$T_{k,y}^m(x) := \sum_{\ell=0}^m \frac{1}{\ell!} D^\ell k(y) (x - y)^\ell.$$

Theorem 1.4. *Under the assumptions of Theorem 1.1 suppose $k \in C^6(S^3)$ and let*

$$M_0^* := \{\theta \in M : a_1(\theta) = 0 \text{ and } a_3(\theta) \neq 0\}.$$

Then there is $\delta > 0$ such that for any $\theta \in M_0^$ there exists a unique C^1 -curve*

$$\{0 < \mu < \delta\} \ni \mu \mapsto (t(\mu), \varphi(\mu, \cdot)) \in ((-\delta, 1 + \delta) \setminus \{0\}) \times C^{2,\alpha}(S^3),$$

such that as $\mu \rightarrow 0$

$$t(\mu) = -\frac{a_3(\theta)}{a_2(\theta)} \mu + O(\mu^{1+\frac{1}{4}}),$$

and $\varphi(\mu, \cdot)$ solves (1.5) for $t = t(\mu)$ and blows up like

$$\|\mathcal{R}_\theta(\varphi(\mu, x)) - z_{\mu,0}(x)\|_{\mathcal{D}^{1,2}(\mathbb{R}^3) \cap C^2(B_1(0))} = O(\mu^2).$$

The curve is unique, in the sense that, if $(t_i, \varphi_i) \in ((-\delta, 1 + \delta) \setminus \{0\}) \times C^{2,\alpha}(S^3)$ blow up at $\theta \in M_0^$ then there is a sequence of positive numbers (μ_i) converging to zero such that $(t_i, \varphi_i) = (t(\mu_i), \varphi(\mu_i, \cdot))$ for all but finitely many $i \in \mathbb{N}$.*

To illustrate our results we give an example. Suppose k_θ is given by

$$k_\theta(x) = 1 + \frac{3x_1^2 - 2x_2^2 - x_3^2}{(1 + |x|^2)^2} + \frac{|x|^4}{(1 + |x|^2)^3} \left(b - \frac{a}{(1 + |x|^2)} - \frac{1 - a}{(1 + |x|^2)^2} \right).$$

Then $1 + tk_\theta(x)$ is strictly positive for all $t \geq 0$, if $b \geq 0$ and $a \leq 3$, and $\Delta k_\theta(0) = 0$ and $\nabla k_\theta(0) = \nabla \Delta k_\theta(0) = 0$. Furthermore,

$$\begin{aligned} a_0(\theta) &= -\frac{\pi^2}{64}(35 - 48b + 5a), \quad a_1(\theta) = 120(b - 1), \\ a_2(\theta) &= 120(b - 1) - 56, \quad a_3(\theta) = 75\left(6b + \frac{7}{8}a - \frac{63}{8}\right). \end{aligned}$$

Our results show: θ is not a blow-up point, if $a_0(\theta) \neq 0$, that is $a \neq -7 + \frac{48}{5}b$, or $a_2(\theta) = 0$, that is $b = \frac{22}{15}$. Moreover, if $a_0(\theta) = 0$ and $a_2(\theta) \neq 0$ then there is a curve of solutions $(t(\mu), \varphi(\mu, \cdot))$ which blow up at θ such that

$$\begin{aligned} t(\mu) &= \frac{b-1}{\frac{22}{15}-b} + O(\mu^{\frac{1}{4}}), \quad \text{if } b \neq 1 \\ t(\mu) &= \frac{30}{56}\mu + O(\mu^{1+\frac{1}{4}}), \quad \text{if } b = 1. \end{aligned}$$

We sketch the strategy of the proofs of our main results and outline the remaining part of the paper. The transformation in (1.3) gives rise to a Hilbert space isomorphism between $H^{1,2}(S^N)$ and $\mathcal{D}^{1,2}(\mathbb{R}^N)$, where $\mathcal{D}^{1,2}(\mathbb{R}^N)$ denotes the closure of $C_c^\infty(\mathbb{R}^N)$ with respect to

$$\|u\|^2 := \int_{\mathbb{R}^N} |\nabla u|^2 = \langle u, u \rangle.$$

Due to elliptic regularity (see [8]) and Harnack's inequality it is enough to find a weak nonnegative solution of (1.1) in $H^{1,2}(S^N)$, or of the equivalent equation. Although we take advantage of both formulations, we mainly consider (1.2). We use a finite dimensional reduction of Melnikov type developed in [1, 2] and find solutions of (1.2) as critical points of $f_t : \mathcal{D}^{1,2}(\mathbb{R}^N) \rightarrow \mathbb{R}$, where

$$f_t(u) := \frac{1}{2} \int_{\mathbb{R}^N} |\nabla u|^2 - \frac{N-2}{2N} \int_{\mathbb{R}^N} (1 + tk(x)) |u|^{\frac{2N}{N-2}}.$$

For $t = 0$ the functional f_0 possesses, as seen above, a $N + 1$ dimensional manifold of critical points Z . To setup the finite dimensional reduction we need to analyze Z and the spectrum of $f_0''(z)$ in detail, which is done for all $N \geq 3$ in Section 2. For the rest of the paper we will only deal with the case $N = 3$. In Section 3 we recall without proof that if $N = 3$ a sequence of solutions to (1.5) can only blow-up in a single point (see [18, 21]) and fit this result into our framework. Section 4 contains the finite dimensional reduction of our problem. In contrast to [2], where the reduction is performed for small t , we show that a finite dimensional reduction of (1.6) for large t is still possible. We end up with a function $\vec{\alpha} : U \rightarrow \mathbb{R}^4$, where $U \subset \mathbb{R} \times Z$, such that the zeros of $\vec{\alpha}(t, \cdot)$ correspond to solutions of (1.6) with large L^∞ norm. We recall that Z is parametrized by μ and y . Now, to construct or to rule out blow-up sequences it is enough to construct or exclude zeros of $\vec{\alpha}(t, \cdot)$ for small μ . To this end we need to expand $\vec{\alpha}$ up to order 5 in μ and to compute derivatives of $\vec{\alpha}$, which is done in Sections 4 and 5. We see that θ can only be a blow-up point if $\nabla k(\theta) = 0$ and $\Delta k_\theta(0) = 0$. In Section 6 we finally obtain under the assumptions of Theorems 1.1- 1.4 that there are (t_i, φ_i) which blow up at θ if and only if

$\nabla k(\theta) = 0$, $\Delta k_\theta(0) = 0$, and there exist positive (μ_i) converging to 0 such that

$$0 = a_0(\theta) + \mu_i(a_1(\theta) + t_i a_2(\theta)) + \mu_i^2 a_3(\theta) + O(t_i \mu_i^{1+\frac{1}{4}} + \mu_i^{2+\frac{1}{4}}).$$

This gives our main results, which are stated and proved in Section 7.

In a subsequent paper [20] we use the above a priori estimates and compute under the assumptions of Theorem 1.1 the Leray-Schauder degree d of the problem (1.5). We show that if $M^* = \emptyset$ and k is a Morse function then,

$$d = -\left(1 - \sum_{\theta \in \text{Crit}(k)^-} (-1)^{\text{ind}(\theta)}\right), \quad \text{where}$$

$$\text{Crit}(k)^- := \{\theta \in S^3 : \nabla k(\theta) = 0 \text{ and } \lim_{\mu \rightarrow 0^+} \text{sgn}(\Delta k_\theta(0) + a_0(\theta)\mu + a_1(\theta)\mu^2) = -1\},$$

generalizing the existence result in [5, 10, 21].

Our approach yields information about blow-up sequences as precise as we want, that is of any order in μ or y . For instance it is possible to compute the term of order μ^6 , which is of interest when $a_3(\theta)$ is zero. But the necessary computations and terms, as may already be seen in the expansion of order 5, are getting rather bulky. In higher dimensions $N \geq 4$ solutions may blow up in more than one point and our method, which still applies with minor changes to $N \geq 4$, will only give information about “one bubble” blow-up.

2. PRELIMINARIES

We define for $\mu > 0$ and $y \in \mathbb{R}^N$ the maps $\mathcal{U}_\mu, \mathcal{T}_y : \mathcal{D}^{1,2}(\mathbb{R}^N) \rightarrow \mathcal{D}^{1,2}(\mathbb{R}^N)$ by

$$\mathcal{U}_\mu(u) := \mu^{-\frac{N-2}{2}} u\left(\frac{\cdot}{\mu}\right) \text{ and } \mathcal{T}_y(u) := u(\cdot - y).$$

With this notation the critical manifold Z is given by

$$Z = \{z_{\mu,y} = \mathcal{T}_y \circ \mathcal{U}_\mu(z_{1,0}) : y \in \mathbb{R}^N, \mu > 0\}.$$

It is easy to check that the dilation $\mathcal{U}_\mu$ and the translation $\mathcal{T}_y$ conserve the norms $\|\cdot\|$ and the L^{2^*} -Norm $\|\cdot\|_{2^*}$, where $2^* := 2N/(N-2)$. Thus for every $\mu > 0$ and $y \in \mathbb{R}^N$

$$\begin{aligned} (\mathcal{U}_\mu)^{-1} &= (\mathcal{U}_\mu)^t = U_{\mu^{-1}}, \quad (\mathcal{T}_y)^{-1} = (\mathcal{T}_y)^t = \mathcal{T}_{-y}, \text{ and} \\ f_0 &= f_0 \circ \mathcal{U}_\mu = f_0 \circ \mathcal{T}_y \end{aligned} \tag{2.1}$$

where $(\cdot)^t$ denotes the adjoint. Twice differentiating the identities for f_0 in (2.1) yields

$$f_0''(v) = (\mathcal{T}_y \circ \mathcal{U}_\mu)^{-1} \circ f_0''(\mathcal{T}_y \circ \mathcal{U}_\mu(v)) \circ (\mathcal{T}_y \circ \mathcal{U}_\mu) \quad \forall v \in \mathcal{D}^{1,2}(\mathbb{R}^N). \tag{2.2}$$

Moreover, we see that $U(\mu, y, z) := \mathcal{T}_y \circ \mathcal{U}_\mu(z)$ maps $(0, \infty) \times \mathbb{R}^N \times Z$ into Z , hence

$$\begin{aligned} \frac{\partial U}{\partial z}(\mu, y, z) &= \mathcal{T}_y \circ \mathcal{U}_\mu : T_z Z \rightarrow T_{\mathcal{T}_y \circ \mathcal{U}_\mu(z)} Z \text{ and} \\ \mathcal{T}_y \circ \mathcal{U}_\mu &: (T_z Z)^\perp \rightarrow (T_{\mathcal{T}_y \circ \mathcal{U}_\mu(z)} Z)^\perp. \end{aligned} \tag{2.3}$$

The tangent space $T_{z_{\mu,y}}Z$ at a point $z_{\mu,y} \in Z$ is spanned by $N+1$ orthonormal functions $\dot{\xi}_{\mu,y}^i$,

$$T_{z_{\mu,y}}Z = \langle \dot{\xi}_{\mu,y}^i : i = 0 \dots N \rangle,$$

where $\dot{\xi}_{\mu,y}^i$ denotes for $i = 0$ the normalized tangent vector $\frac{d}{d\mu}z_{\mu,y}$ and for $1 \leq i \leq N$ the normalized tangent vector $\frac{d}{dy_i}z_{\mu,y} = -\frac{\partial}{\partial x_i}z_{\mu,y}$. By (2.3) we obtain

$$\dot{\xi}_{\mu,y}^i = \mathcal{T}_y \circ \mathcal{U}_\mu(\dot{\xi}_{1,0}^i).$$

An explicit calculation gives for $1 \leq i \leq N$

$$(\dot{\xi}_{1,0}^i)_i = \tau_1(1 + |x|^2)^{-\frac{N}{2}}x_i, \quad \tau_1^2 := \frac{\Gamma(N+2)}{\pi^{N/2}\Gamma(2+N/2)N}.$$

For $i = 0$ we find

$$(\dot{\xi}_{1,0}^0)_0 = \tau_0(1 + |x|^2)^{-\frac{N-2}{2}}\left(1 - \frac{2}{(1 + |x|^2)}\right), \quad \tau_0^2 := \frac{(N+1)\Gamma(N)}{\pi^{N/2}\Gamma(N/2)N(N+2)}.$$

Using the canonical identification of the Hilbert space $\mathcal{D}^{1,2}(\mathbb{R}^N)$ with its dual induced by the scalar-product we shall consider $f'_t(u)$ as an element of $\mathcal{D}^{1,2}(\mathbb{R}^N)$ and $f''_t(u)$ as one of $\mathcal{L}(\mathcal{D}^{1,2}(\mathbb{R}^N))$. With this identification $f''_t(u)$ is of the form *identity - compact* (see [2]) and hence a Fredholm operator of index zero.

Since $f''_0(z_{\mu,y})$ is a self-adjoint, compact perturbation of the identity map in $\mathcal{D}^{1,2}(\mathbb{R}^N)$, its spectrum $\sigma(f''_0(z_{\mu,y}))$ consists of point-spectrum, possibly accumulating at 1. We fix $\lambda \in \sigma(f''_0(z_{\mu,y}))$ and a corresponding eigenfunction u . Then u solves

$$-\Delta u - \frac{N+2}{N-2}z_{\mu,y}^{2^*-2}u = \lambda(-\Delta u). \quad (2.4)$$

We expand u in spherical harmonics with center y

$$u(y + r\vartheta) = \sum_{i=0}^{\infty} \sum_{l=1}^{c_i} v_{i,l}(r)Y_{i,l}(\vartheta), \quad r \in \mathbb{R}^+, \quad \vartheta \in \mathbb{S}^{N-1},$$

where

$$v_{i,l}(r) = \int_{\mathbb{S}^{N-1}} u(y + r\vartheta)Y_{i,l}(\vartheta) d\vartheta, \quad c_i := \binom{N-1+i}{N-1} - \binom{N-3+i}{N-1},$$

and $\{Y_{i,l}\}$ denote a $L^2(\mathbb{S}^{N-1})$ -basis of (real valued) spherical harmonics satisfying for all $i \in \mathbb{N}_0$ and $1 \leq l \leq c_i$

$$-\Delta_{\mathbb{S}^{N-1}}Y_{i,l} = i(N+i-2)Y_{i,l}.$$

There is a freedom in choosing such a L^2 -basis and because the cases $i = 1, 2$ will be of special interest in the sequel we fix the basis-vectors in these cases. We set for $i = 1$ and $1 \leq l \leq N$

$$Y_{1,l}\left(\frac{x}{|x|}\right) := \left(\frac{2\pi^{N/2}}{N\Gamma(N/2)}\right)^{-1/2} \frac{x_l}{|x|}. \quad (2.5)$$

For $i = 2$ we introduce a more convenient notation and write

$$Y_{2,(l_1,l_2)}\left(\frac{x}{|x|}\right) := \left(\frac{\pi^{N/2}}{\Gamma(2+N/2)}\right)^{-1/2} |x|^{-2} \\ \times \begin{cases} \sqrt{2} x_{l_1} x_{l_2} & \text{if } 1 \leq l_1 < l_2 \leq N \\ \sqrt{\frac{l_1-1}{l_1}} \left(x_{l_1}^2 - \frac{1}{l_1-1} \sum_{m=1}^{l_1-1} x_m^2\right) & \text{if } 2 \leq l_1 = l_2 \leq N. \end{cases} \quad (2.6)$$

Since u solves (2.4) the functions $v_{i,l}$ satisfy for $i \in \mathbb{N}_0$ and $1 \leq l \leq c_i$

$$-v_{i,l}'' - \frac{N-1}{r} v_{i,l}' + \frac{i(N+i-2)}{r^2} v_{i,l} = \frac{N+2}{(N-2)(1-\lambda)} z_{\mu,0}^{2*-2} v_{i,l}.$$

Making the transformation

$$v(r) = r^{-\frac{N-2}{2}} \zeta(\ln \mu + \ln r),$$

we obtain the equation for $i \in \mathbb{N}_0$ and $1 \leq l \leq c_i$

$$-\zeta_{i,l}'' - \frac{N(N+2)}{4(1-\lambda)} \cosh^{-2}(t) \zeta_{i,l} = \left(-\left(\frac{N-2}{2}\right)^2 - i(N+i-2)\right) \zeta_{i,l}.$$

Using the results in [17, p. 74] or [15] as in [13] we find

$$\lambda_{i,j} = 1 - \frac{N(N+2)}{((N+1) + 2(i+j-1))^2 - 1}.$$

The corresponding eigenfunction is given by

$$\psi_{i,j}(t) := (1 - \tanh(t)^2)^{\frac{N-2+2i}{4}} \mathcal{P}_j^{(\sigma_i, \sigma_i)}(\tanh(t)),$$

where $\mathcal{P}_j^{(\sigma, \sigma)}$ denotes the *Jacobi* polynomial defined in (A.1) and σ_i is given by

$$\sigma_i := \frac{N-2}{2} + i.$$

Consequently, $\sigma(f_0''(z_{\mu,y})) = \{\lambda_{i,j} : i, j \in \mathbb{N}_0\}$ and the eigenspace of the eigenvalue $\lambda_{i,j}$ has dimension c_i and is spanned by, ($l = 1 \dots c_i$)

$$\Phi_{i,j,l}^{\mu,y}(x) := a_{i,j} \mathcal{T}_y \circ \mathcal{U}_\mu \left(|x|^i (1 + |x|^2)^{-\frac{N-2}{2}-i} \right. \\ \left. \mathcal{P}_j^{(\sigma_i, \sigma_i)}(1 - 2(1 + |x|^2)^{-1}) Y_{i,l}\left(\frac{x}{|x|}\right) \right), \quad (2.7)$$

where the $a_{i,j}$ are given by

$$a_{i,j}^2 := \frac{2(N-1+2(i+j))j!\Gamma(N-1+2i+j)}{(N-2+2(i+j))(N+2(i+j))\Gamma(N/2+i+j)^2}$$

to assure that the $\Phi_{i,j,l}^{\mu,y}$ are orthonormal. Since Z is a manifold of critical points of f_0' , the tangent space $T_z Z$ at a point $z \in Z$ is contained in the kernel $N(f_0''(z))$ of $f_0''(z)$. As $\lambda_{i,j} = 0$ if and only if $i+j = 1$, the dimension of $N(f_0''(z))$ is $N+1$, which implies that

$$T_z Z = N(f_0''(z)) \quad \text{for all } z \in Z. \quad (2.8)$$

More precisely, we have

$$\Phi_{1,0,l}^{\mu,y} = \xi_{\mu,y}^l \text{ for } l = 1, \dots, N \text{ and } \Phi_{0,1,1}^{\mu,y} = \xi_{\mu,y}^0. \quad (2.9)$$

If (2.8) holds the critical manifold Z is called non-degenerate (see [1]) and the self-adjoint Fredholm operator $f_0''(z)$ maps the space $\mathcal{D}^{1,2}(\mathbb{R}^N)$ into $T_z Z^\perp$ and is invertible in $\mathcal{L}(T_z Z^\perp)$. From (2.2) and (2.3), we obtain in this case

$$\|(f_0''(z_{1,0}))^{-1}\|_{\mathcal{L}(T_{z_{1,0}} Z^\perp)} = \|(f_0''(z))^{-1}\|_{\mathcal{L}(T_z Z^\perp)} \quad \forall z \in Z. \quad (2.10)$$

3. BLOW UP ANALYSIS

Let $(K_i) \in C^1(S^3)$ satisfy for some A_0

$$A_0^{-1} \leq K_i(x) \leq A_0 \text{ and } \|\nabla K_i\|_\infty \leq A_0. \quad (3.1)$$

We have the following result (see [18, 21])

Theorem 3.1. *Suppose $(K_i) \in C^1(S^3)$ satisfies (3.1) and $(\varphi_i) \in C^2(S^3)$ solves (1.1) with $N = 3$. Then after passing to a subsequence either (φ_i) is uniformly bounded in $L^\infty(S^3)$ and hence in $C^{2,\alpha}(S^3)$ by elliptic regularity or (φ_i) has precisely one isolated simple blow-up point θ , i.e. there exists a sequence (θ_i) of maxima of φ_i converging to some $\theta \in S^3$ and $C = C(A_0)$ such that $\varphi_i(\theta_i) \rightarrow +\infty$ and in geodesic normal coordinates about θ_i given by $\exp_{\theta_i}(\cdot)$*

$$i\varphi_i(\theta_i)^{-2} \rightarrow 0,$$

$$\left\| \frac{\varphi_i\left(\exp_{\theta_i}\left(\frac{x}{\varphi_i(\theta_i)^2}\right)\right)}{\varphi_i(\theta_i)} - \left(1 + \frac{K_i(\theta_i)}{24}|x|^2\right)^{-\frac{1}{2}} \right\|_{C^{2,\alpha}(\{|x| \leq 3i\})} \leq i^{-4},$$

$$\varphi_i(x) \leq C\varphi_i(\theta_i)^{-1} \text{dist}_{S^3}(x, \theta_i)^{-1} \text{ for } \text{dist}_{S^3}(x, \theta_i) \geq i\varphi_i(\theta_i)^{-2}.$$

We need a slightly different version of this result.

Corollary 3.2. *Under the assumptions of Theorem 3.1 the sequence $(\varphi_i) \in C^2(S^3)$ is, after passing to a subsequence, either uniformly bounded in $C^{2,\alpha}$ or there exist $\theta \in S^3$ and sequences $(\mu_i) \in (0, \infty)$, $(y_i) \in \mathbb{R}^3$ satisfying*

$$\lim_{i \rightarrow \infty} \mu_i = 0, \quad \lim_{i \rightarrow \infty} y_i = 0,$$

such that in stereographic coordinates $\mathcal{S}_\theta(\cdot)$ about θ the function u_i defined by the transformation (1.3) satisfies

$$u_i - 6^{\frac{1}{4}}(K_i \circ \mathcal{S}_\theta(y_i))^{-\frac{1}{4}} z_{\mu_i, y_i} \text{ is orthogonal to } T_{z_{\mu_i, y_i}} Z, \quad (3.2)$$

$$\|u_i - 6^{\frac{1}{4}}(K_i \circ \mathcal{S}_\theta(y_i))^{-\frac{1}{4}} z_{\mu_i, y_i}\|_{\mathcal{D}^{1,2}(\mathbb{R}^3)} = o_{A_0}(1). \quad (3.3)$$

To prove the corollary we first need the following lemma, which is an easy consequence of Theorem 3.1.

Lemma 3.3. *Under the assumptions of Theorem 3.1 the sequence $(\varphi_i) \in C^2(S^3)$ is, after passing to a subsequence, either uniformly bounded in $C^{2,\alpha}$ or there exists a sequence (θ_i) of maxima of φ_i converging to some $\theta \in S^3$ and such that $\varphi_i(\theta_i) \rightarrow +\infty$ and in stereographic coordinates $\mathcal{S}_\theta(\cdot)$ using the transformation (1.3)*

$$\|u_i - 6^{\frac{1}{4}} K_i(\theta_i)^{-\frac{1}{4}} z_{\mu_i, y_i}(x)\|_{\mathcal{D}^{1,2}(\mathbb{R}^3)} = o_{A_0}(1),$$

where $\mu_i \rightarrow 0$ and $y_i \rightarrow 0$ are given by

$$y_i := \mathcal{S}_\theta^{-1}(\theta_i), \quad \mu_i := (6/K_i(\theta_i))^{\frac{1}{2}} \varphi_i(\theta_i)^{-2}.$$

Proof of Corollary 3.2. From Lemma 3.3 we infer that there are $\theta \in S^3$, $(\tilde{\mu}_i)$, and $(\tilde{y}_i)$ such that

$$\mu_i + |y_i| + \|u_i - 6^{\frac{1}{4}} (K_i \circ \mathcal{S}_\theta(\tilde{y}_i))^{-\frac{1}{4}} z_{\tilde{\mu}_i, \tilde{y}_i}\|_{\mathcal{D}^{1,2}(\mathbb{R}^3)} = o_{A_0}(1).$$

For i fixed, consider

$$d_i := \inf_{\mu, y} \|u_i - 6^{\frac{1}{4}} (K_i \circ \mathcal{S}_\theta(\tilde{y}_i))^{-\frac{1}{4}} z_{\mu, y}\|_{\mathcal{D}^{1,2}(\mathbb{R}^3)}^2.$$

Clearly $d_i = o_{A_0}(1)$ and therefore d_i is attained at μ_i, y_i and

$$u_i - 6^{\frac{1}{4}} (K_i \circ \mathcal{S}_\theta(\tilde{y}_i))^{-\frac{1}{4}} z_{\mu_i, y_i} \text{ is orthogonal to } T_{z_{\mu_i, y_i}} Z.$$

Since z_{μ_i, y_i} is orthogonal to $T_{z_{\mu_i, y_i}} Z$ relation (3.2) follows. To prove rest of the claim we need to estimate $|y_i - \tilde{y}_i|$ and $|\mu_i - \tilde{\mu}_i|$. To this end we observe that by construction

$$o_{A_0}(1) = \|z_{\mu_i, y_i} - z_{\tilde{\mu}_i, \tilde{y}_i}\|_{\mathcal{D}^{1,2}(\mathbb{R}^3)} = \|z_{\tilde{\mu}_i/\mu_i, \tilde{y}_i - y_i} - z_{1,0}\|_{\mathcal{D}^{1,2}(\mathbb{R}^3)}.$$

Since

$$\lim_{\mu + \mu^{-1} + |y| \rightarrow \infty} \|z_{\mu, y} - z_{1,0}\|_{\mathcal{D}^{1,2}(\mathbb{R}^3)}^2 = 2\|z_{1,0}\|_{\mathcal{D}^{1,2}(\mathbb{R}^3)}^2$$

we see that there is $R_1 = R_1(A_0) > 0$ such that

$$(R_1)^{-1} \leq \tilde{\mu}_i/\mu_i \leq R_1 \text{ and } |y_i - \tilde{y}_i| \leq R_1.$$

Now by explicit calculations or elliptic regularity [8] we have

$$\max \left\{ |y_i - \tilde{y}_i|, \left| \frac{\tilde{\mu}_i}{\mu_i} - 1 \right| \right\} \leq \text{const}(A_0) \|z_{\tilde{\mu}_i/\mu_i, \tilde{y}_i - y_i} - z_{1,0}\|_{\mathcal{D}^{1,2}(\mathbb{R}^3)} = o_{A_0}(1),$$

which gives the claim. $\square$

4. EXPANSION OF THE PERTURBATION TERMS w AND $\vec{\alpha}$

For the rest of the paper we will only treat the case $N = 3$. Unless otherwise indicated, integration extends over $\mathbb{R}^3$ and is done with respect to the variable x . Moreover, we will write k instead of k_θ when there is no possibility of confusion to avoid cumbersome subindexing.

From the change of coordinates $x \mapsto \mu x + y$, Hölder's and Sobolev's inequality we get

Lemma 4.1. *Let $y \in \mathbb{R}^3$, $\tau > 0$ and $f, r : \mathbb{R}^3 \rightarrow \mathbb{R}$ measurable such that*

$$|r(x)| \leq C_r |x - y|^\sigma \text{ in } B_1(y), \quad |r(x)| \leq C_r |x - y|^{\tilde{\sigma}} \text{ in } \mathbb{R}^3 \setminus B_1(y),$$

$$|f(x)| \leq C_r |x|^{-s} \text{ in } B_1(0), \quad |f(x)| \leq C_r |x|^{-m} \text{ in } \mathbb{R}^3 \setminus B_1(0),$$

for some $C_r, m, s > 0$ and $0 \leq \tilde{\sigma}, \sigma$. Then there is $C = C(\tau, C_r) > 0$ such that for $v, v_1, v_2 \in \mathcal{D}^{1,2}(\mathbb{R}^3)$:

For $0 \leq \tilde{\sigma}, \sigma \leq m - 3 - \tau$ and $s + \tau \leq 3 + \sigma$ there holds

$$\left| \int r(x) \mu^{-3} f\left(\frac{x - y}{\mu}\right) \right| \leq C (\chi_{(0,1]}(\mu) \mu^\sigma + \chi_{(1,\infty)}(\mu) (\mu^{s-3} + \mu^{\tilde{\sigma}})),$$

if $0 \leq \tilde{\sigma}, \sigma \leq m - \frac{5}{2} - \tau$ and $s + \tau < \sigma + \frac{5}{2}$ then

$$\left\| \int \frac{r(x)}{\mu^{\frac{5}{2}}} f\left(\frac{x-y}{\mu}\right) \cdot \right\|_{\mathcal{D}^{1,2}(\mathbb{R}^3)} \leq C(\chi_{(0,1]}(\mu)\mu^\sigma + \chi_{(1,\infty)}(\mu)(\mu^{s-\frac{5}{2}} + \mu^{\tilde{\sigma}})),$$

if $0 \leq \tilde{\sigma}, \sigma \leq m - 2 - \tau$ and $s + \tau < \sigma + 2$ then

$$\begin{aligned} \sup_{\|v\| \leq 1} \left\| \int r(x)\mu^{-2} f\left(\frac{x-y}{\mu}\right) v \cdot \right\|_{\mathcal{D}^{1,2}(\mathbb{R}^3)} \\ \leq C(\chi_{(0,1]}(\mu)\mu^\sigma + \chi_{(1,\infty)}(\mu)(\mu^{s-2} + \mu^{\tilde{\sigma}})), \end{aligned}$$

if $0 \leq \tilde{\sigma}, \sigma \leq m - \frac{3}{2} - \tau$ and $s + \tau < \sigma + \frac{3}{2}$ then we have

$$\begin{aligned} \sup_{\|v_1\|, \|v_2\| \leq 1} \left\| \int r(x)\mu^{-\frac{3}{2}} f\left(\frac{x-y}{\mu}\right) v_1 v_2 \cdot \right\|_{\mathcal{D}^{1,2}(\mathbb{R}^3)} \\ \leq C(\chi_{(0,1]}(\mu)\mu^\sigma + \chi_{(1,\infty)}(\mu)(\mu^{s-\frac{3}{2}} + \mu^{\tilde{\sigma}})). \end{aligned}$$

Using the above estimates we may prove the main ingredient for the finite dimensional reduction.

Lemma 4.2. Suppose $k \in C^5(\mathbb{R}^3)$ and there are $A_0, B_0, B_1 > 0$ such that

$$\max(B_0, B_1, \sup_{|m| \leq 5} \|D^m k\|_\infty) \leq A_0 \text{ and}$$

$$A_0^{-1} \leq 1 + tk(x) \quad \forall (x, t) \in \mathbb{R}^3 \times [-B_0, B_1].$$

Then there exist $\rho_0 = \rho_0(A_0) > 0$, $t_0 = t_0(A_0) > 0$, an upper continuous function $\mu_0 : \mathbb{R} \rightarrow \mathbb{R}_+ \cup \{\infty\}$, depending only on A_0 , and two functions $w : \Omega \rightarrow \mathcal{D}^{1,2}(\mathbb{R}^3)$ and $\vec{\alpha} : \Omega \rightarrow \mathbb{R}^4$, where

$$\begin{aligned} \Omega &:= \{(t, \mu, y) \in [-B_0, B_1] \times (0, +\infty) \times \mathbb{R}^3 : 0 < \mu < \mu_0(t)\}, \\ \mu_0(t) &= +\infty \text{ if } |t| \leq t_0, \end{aligned}$$

such that for any $(t, \mu, y) \in \Omega$

$$w(t, \mu, y) \text{ is orthogonal to } T_{z_{\mu,y}} Z \quad (4.1)$$

$$f'_t(z_{\mu,y} + w(t, \mu, y)) = \vec{\alpha}(t, \mu, y) \cdot \dot{\xi}_{\mu,y} \in T_{z_{\mu,y}} Z \quad (4.2)$$

$$\|w(t, \mu, y) - w_0(t, \mu, y)\| + \|\vec{\alpha}(t, \mu, y)\| < \rho_0, \quad (4.3)$$

where $\{\dot{\xi}_{\mu,y}^i : i = 0 \dots 3\}$ denotes the orthonormal basis of $T_{z_{\mu,y}} Z$ given in (2.9) and

$$w_0(t, \mu, y) := ((1 + tk(y))^{-\frac{1}{4}} - 1)z_{\mu,y}.$$

The functions w and $\vec{\alpha}$ are of class C^2 and unique in the sense that if $(v, \vec{\beta})$ satisfies (4.1)-(4.3) for some $(t, \mu, y) \in \Omega$ then $(v, \vec{\beta})$ is given by $(w(t, \mu, y), \vec{\alpha}(t, \mu, y))$. Moreover, we have for $1 \leq j \leq 3$

$$\begin{aligned} \|w(t, \mu, y) - \sum_{i=0}^2 w_i(t, \mu, y)\| + \|\vec{\alpha}(t, \mu, y)_0 - \sum_{i=1}^2 \vec{\alpha}_i(t, \mu, y)_0\| \\ \leq O_{A_0} \left(t(|\nabla k(y)|^2 \min(1, \mu^2) + \min(1, \mu^{\frac{9}{4}})) \right), \end{aligned} \quad (4.4)$$

$$\|\vec{\alpha}(t, \mu, y)_j - \sum_{i=1}^2 \vec{\alpha}_i(t, \mu, y)_j\| \leq O_{A_0} \left(t \min(1, \mu^{\frac{9}{4}}) \right),$$

where

$$\begin{aligned}\vec{\alpha}_1(t, \mu, y) &:= -t \min(1, \mu)(1 + tk(y))^{-\frac{5}{4}} \frac{\pi}{3^{\frac{1}{4}}\sqrt{5}} \begin{pmatrix} 0 \\ \nabla k(y) \end{pmatrix}, \\ \vec{\alpha}_2(t, \mu, y) &:= -t \min(1, \mu^2)(1 + tk(y))^{-\frac{5}{4}} \frac{\pi}{3^{\frac{1}{4}}\sqrt{5}} \begin{pmatrix} \Delta k(y) \\ \vec{0} \end{pmatrix}.\end{aligned}$$

and

$$\begin{aligned}w_1(t, \mu, y) &:= t \min(1, \mu)(1 + tk(y))^{-\frac{5}{4}} \mathcal{T}_y \circ \mathcal{U}_\mu(\tilde{w}_1(y)), \\ \tilde{w}_1(y) &:= \mathcal{F}_0^{-1} \left(\int \nabla k(y) x(z_{1,0})^5 \cdot \right), \\ w_2(t, \mu, y) &:= t \min(1, \mu^2)(1 + tk(y))^{-\frac{5}{4}} \mathcal{T}_y \circ \mathcal{U}_\mu(\tilde{w}_2(y)), \\ \tilde{w}_2(y) &:= \mathcal{F}_0^{-1} \left(\frac{1}{2} \int D^2 k(y) x^2(z_{1,0})^5 \cdot \right),\end{aligned}$$

The operator $\mathcal{F}_0^{-1} \in \mathcal{L}(\mathcal{D}^{1,2}(\mathbb{R}^3), T_{z_{1,0}} Z^\perp)$ is defined by

$$\mathcal{F}_0^{-1} := (f_0''(z_{1,0})|_{T_{z_{1,0}} Z^\perp})^{-1} \circ \text{Proj}_{T_{z_{1,0}} Z^\perp}.$$

Proof. Define $H : \mathbb{R} \times (0, \infty) \times \mathbb{R}^3 \times \mathcal{D}^{1,2}(\mathbb{R}^3) \times \mathbb{R}^4 \rightarrow \mathcal{D}^{1,2}(\mathbb{R}^3) \times \mathbb{R}^4$

$$H(t, \mu, y, w, \vec{\alpha}) := (f_t'(z_{\mu,y} + w) - \vec{\alpha} \cdot \dot{\xi}_{\mu,y}, (\langle w, (\dot{\xi}_{\mu,y})_l \rangle)_l).$$

If $H(t, \mu, y, w, \vec{\alpha}) = (0, 0)$ then w satisfies (4.1)-(4.2). We have

$$\left(\frac{\partial H}{\partial(w, \vec{\alpha})}(t, \mu, y, w, \vec{\alpha}) \right) \begin{pmatrix} \varphi \\ \vec{\beta} \end{pmatrix} = (f_t''(z_{\mu,y} + w)\varphi - \vec{\beta} \dot{\xi}_{\mu,y}, (\langle \varphi, (\dot{\xi}_{\mu,y})_l \rangle)_l). \quad (4.5)$$

Note that

$$\begin{aligned}\left\langle \left(\frac{\partial H}{\partial(w, \vec{\alpha})}(0, \mu, y, 0, 0) \right) \begin{pmatrix} w \\ \vec{\beta} \end{pmatrix}, (f_0''(z_{\mu,y})w - \vec{\beta} \cdot \dot{\xi}_{\mu,y}, (\langle w, (\dot{\xi}_{\mu,y})_l \rangle)_l) \right\rangle \\ = \|f_0''(z_{\mu,y})w\|^2 + |\vec{\beta}|^2 + |(w, \dot{\xi}_{\mu,y})_i|^2.\end{aligned} \quad (4.6)$$

From (2.10) and (4.6) we infer that $(\frac{\partial H}{\partial(w, \vec{\alpha})}(0, \mu, y, 0, 0))$ is an injective Fredholm operator of index zero, hence invertible and

$$\left\| \left(\frac{\partial H}{\partial(w, \vec{\alpha})}(0, \mu, y, 0, 0) \right)^{-1} \right\| \leq 1 + \|(f_0''(z_{\mu,y}))^{-1}\|_{\mathcal{L}(T_{z_{\mu,y}} Z^\perp)} =: C_*. \quad (4.7)$$

Clearly, $H(t, \mu, y, w, \vec{\alpha}) = (0, 0)$ if and only if $(w, \vec{\alpha}) = F_{t,\mu,y}(w, \vec{\alpha})$, where

$$F_{t,\mu,y}(w, \vec{\alpha}) := - \left(\frac{\partial H}{\partial(w, \vec{\alpha})}(0, \mu, y, 0, 0) \right)^{-1} H(t, \mu, y, w, \vec{\alpha}) + (w, \vec{\alpha}).$$

We will prove that $F_{t,\mu,y}(w, \vec{\alpha})$ is a contraction in some ball

$$B_\rho \left(\sum_{i=0}^2 w_i(t, \mu, y), \sum_{i=1}^2 \vec{\alpha}_i(t, \mu, y) \right)$$

for any radius ρ such that

$$O_{A_0} \left(t(|\nabla k(y)| \min(1, \mu^2) + \min(1, \mu^{2+\frac{1}{4}})) \right) \leq \rho \leq \rho_0,$$

where $\rho_0 = \rho_0(A_0)$ will be chosen later.

To this end we fix $\rho > 0$ and $(w, \vec{\alpha}) \in B_\rho(0, 0)$. In the sequel we will suppress the dependence of $\vec{\alpha}_i$ and w_i on t, μ and y . From (4.7) and Sobolev's inequality

$$\begin{aligned}
& \frac{1}{C_*} \|F_{t,\mu,y}(w + \sum_{i=0}^2 w_i, \vec{\alpha} + \sum_{i=1}^2 \vec{\alpha}_i) - (\sum_{i=0}^2 w_i, \sum_{i=1}^2 \vec{\alpha}_i)\| \\
& \leq \|f'_t(z_{\mu,y} + w + \sum_{i=0}^2 w_i) - \sum_{i=1}^2 \vec{\alpha}_i \cdot \dot{\xi}_{\mu,y} - f''_0(z_{\mu,y})w\| \\
& \leq \|\langle z_{\mu,y} + \sum_{i=0}^2 w_i - \sum_{i=1}^2 \vec{\alpha}_i \cdot \dot{\xi}_{\mu,y}, \cdot \rangle - 5 \int (z_{\mu,y})^4 w \cdot \\
& \quad - \int (1 + tk(x)) \left(\sum_{l=0}^2 \binom{5}{l} (z_{\mu,y} + w_0)^{5-l} (w_1 + w_2 + w)^l \right) \cdot \| \\
& \quad + O_{A_0}(\|w_1 + w_2 + w\|^3). \tag{4.8}
\end{aligned}$$

Obviously, $(1 + tk(y))(z_{\mu,y} + w_0)^4 = (z_{\mu,y})^4$ and

$$\langle z_{\mu,y} + w_0, \varphi \rangle = \int (1 + tk(y))(z_{\mu,y} + w_0)^5 \varphi. \tag{4.9}$$

Inserting this in (4.8) and using Lemma 4.1 we get

$$\begin{aligned}
& \frac{1}{C_*} \|F_{t,\mu,y}(w + \sum_{i=0}^2 w_i, \vec{\alpha} + \sum_{i=1}^2 \vec{\alpha}_i) - (\sum_{i=0}^2 w_i, \sum_{i=1}^2 \vec{\alpha}_i)\| \\
& \leq \|f''_0(z_{\mu,y})(w_1 + w_2) - \langle (\vec{\alpha}_1 + \vec{\alpha}_2) \cdot \dot{\xi}_{\mu,y}, \cdot \rangle \\
& \quad - t \int (k(x) - k(y))(z_{\mu,y} + w_0)^5 \cdot \| \\
& \quad + O_{A_0}\left(t^2(|\nabla k(y)| \min(1, \mu^2) + \min(1, \mu^3))\right) \\
& \quad + O_{A_0}\left(t \min(1, \mu) \|w\| + \|w\|^2\right). \tag{4.10}
\end{aligned}$$

From (2.1)-(2.3) and the definition of w_1 and w_2 we infer

$$\begin{aligned}
f''_0(z_{\mu,y})w_1 &= t \min(1, \mu^{-1}) \text{Proj}_{T_{z_{\mu,y}}Z^\perp} \left(\int \nabla k(y)(x - y)(z_{\mu,y} + w_0)^5 \cdot \right), \\
f''_0(z_{\mu,y})w_2 &= t \min(1, \mu^{-2}) \text{Proj}_{T_{z_{\mu,y}}Z^\perp} \left(\int \frac{1}{2} D^2 k(y)(x - y)^2 (z_{\mu,y} + w_0)^5 \cdot \right).
\end{aligned}$$

To find $\vec{\alpha}_1$ and $\vec{\alpha}_2$ we observe that since $(\dot{\xi}_{\mu,y})_0$ is even and $(\dot{\xi}_{\mu,y})_i$ is odd for $1 \leq i \leq 3$ we have

$$\begin{aligned}
& \int \nabla k(y)(x - y)(z_{\mu,y} + w_0)^5 (\dot{\xi}_{\mu,y})_0 = 0, \\
& \int \frac{1}{2} D^2 k(y)(x - y)^2 (z_{\mu,y} + w_0)^5 (\dot{\xi}_{\mu,y})_i = 0 \text{ for } 1 \leq i \leq 3.
\end{aligned}$$

For $1 \leq i \leq 3$ we get from (A.4)- (A.7)

$$\int \nabla k(y)(x-y)(z_{\mu,y} + w_0)^5 (\dot{\xi}_{\mu,y})_i = \mu \frac{\pi}{3^{\frac{1}{4}}\sqrt{5}} (1 + tk(y))^{-\frac{5}{4}} \frac{\partial k}{\partial x_i}(y).$$

For $i = 0$ we get

$$\int \frac{1}{2} D^2 k(y)(x-y)^2 (z_{\mu,y} + w_0)^5 (\dot{\xi}_{\mu,y})_i = \mu^2 \frac{\pi}{3^{\frac{1}{4}}\sqrt{5}} (1 + tk(y))^{-\frac{5}{4}} \Delta k(y).$$

Finally, we obtain

$$\begin{aligned} & f_0''(z_{\mu,y})(w_1 + w_2) - (\vec{\alpha}_1 + \vec{\alpha}_2) \cdot \dot{\xi}_{\mu,y} \\ &= t \left(\int \sum_{\ell=1}^2 \frac{D^\ell k(y)}{\ell!} (x-y)^\ell (z_{\mu,y} + w_0)^5 \cdot \right) + O_{A_0}(t\chi_{(1,\infty)}(\mu)), \end{aligned} \quad (4.11)$$

which implies together with (4.10) and Lemma 4.1

$$\begin{aligned} & \frac{1}{C_*} \|F_{t,\mu,y}(w + \sum_{i=0}^2 w_i, \vec{\alpha} + \sum_{i=1}^2 \vec{\alpha}_i) - (\sum_{i=0}^2 w_i, \sum_{i=1}^2 \vec{\alpha}_i)\| \\ & \leq O_{A_0} \left(t(|\nabla k(y)| \min(1, \mu^2) + \min(1, \mu^{2+\frac{1}{4}})) \right) \\ & \quad + O_{A_0} \left(t \min(1, \mu) \|w\| + \|w\|^2 \right). \end{aligned} \quad (4.12)$$

Consequently, if we fix $0 < \rho_0 < 1/4$ we obtain functions μ_0 and t_0 depending on ρ_0 and A_0 such that $F_{t,\mu,y}$ maps $B_\rho(\sum_{i=0}^2 w_i, \sum_{i=1}^2 \vec{\alpha}_i)$ into itself for every $(t, \mu, y) \in \Omega$ and $\rho > 0$ satisfying

$$\text{const}(A_0, \rho_0) |t| (|\nabla k(y)| \min(1, \mu^2) + \min(1, \mu^{2+\frac{1}{4}})) < \rho \leq \rho_0. \quad (4.13)$$

To show that $F_{t,\mu,y}$ is a contraction we fix $\rho > 0$ and two vectors $(v_1, \vec{\beta}_1)$ and $(v_2, \vec{\beta}_2)$ in $B_\rho(0, 0)$. Then using Lemma 4.1 and (4.7)

$$\begin{aligned} & \frac{\|F_{t,\mu,y}(\sum_{i=0}^2 w_i + v_1, \sum_{i=1}^2 \vec{\alpha}_i + \vec{\beta}_1) - F_{t,\mu,y}(\sum_{i=0}^2 w_i + v_2, \sum_{i=1}^2 \vec{\alpha}_i + \vec{\beta}_2)\|}{C_* \|(v_1, \beta_1) - (v_2, \beta_2)\|} \\ & \leq \int_0^1 \|f_t''(z_{\mu,y} + \sum_{i=0}^2 w_i + v_1 + s(v_2 - v_1)) - f_0''(z_{\mu,y})\| ds \\ & \leq \int_0^1 \|f_t''(z_{\mu,y} + w_0) - f_0''(z_{\mu,y})\| ds + O_{A_0}(\rho + \|w_1 + w_2\|) \\ & \leq O_{A_0}(\rho + t \min(1, \mu)). \end{aligned}$$

Thus we have shown that there are $\rho_0 > 0$ and two function μ_0 and t_0 depending only on A_0 , as claimed above, such that $F_{t,\mu,y}$ is a contraction in $B_\rho(\sum_{i=0}^2 w_i, \sum_{i=1}^2 \vec{\alpha}_i)$ for every $(t, \mu, y) \in \Omega$ and $\rho > 0$ satisfying (4.13). From Banach's fixed-point theorem we deduce the existence and uniqueness of the functions w and $\vec{\alpha}$. The usual inverse function theorem yields the C^2 dependence. The estimates in (4.4) hold due to the uniqueness of the fixed-point and because $F_{t,\mu,y}$ is a contraction for every $\rho > 0$ satisfying (4.13). $\square$

We need a precise expansion of w and $\vec{\alpha}$ in critical points y of k in terms of μ and t . This will be done up to order 5 in μ . We later see that we may assume $|\nabla k(y)|$ to be of order $O(\mu^2)$. First we compute $\tilde{w}_2(y)$ in terms of the eigenfunctions of $f_0''(z_{1,0})$.

Lemma 4.3. *Under the assumptions of Lemma 4.2 we have*

$$\begin{aligned} \tilde{w}_2(y) = & \frac{3^{\frac{5}{4}}\sqrt{\pi}}{8} \sum_{j=0}^{\infty} \frac{\Gamma(j + \frac{9}{2})(5 + 2j)a_{2,j}}{\Gamma(j + 6)} \\ & \times \left(\sum_{1 \leq l < m \leq 3} \psi_{l,m}(y) \Phi_{2,j,(l,m)}^{1,0} + \sum_{l=2}^3 \psi_{l,l}(y) \Phi_{2,j,(l,l)}^{1,0} \right) \\ & - \frac{\pi 3^{\frac{3}{4}}}{16} \Delta k(y) \Phi_{0,0,1}^{1,0} + \frac{3^{\frac{1}{4}}\pi}{4} \Delta k(y) \sum_{j=2}^{\infty} \frac{\Gamma(j + \frac{5}{2})(1 + 2j)a_{0,j}}{\Gamma(j + 2)(j + 3)(j - 1)} \Phi_{0,j,1}^{1,0}, \end{aligned}$$

where we use the basis defined in (2.6), (2.7) and

$$\begin{aligned} \psi_{l,m}(y) &:= \frac{2\sqrt{\pi}}{\sqrt{15}} \frac{\partial^2 k(y)}{\partial x_l \partial x_m}, & \psi_{2,2}(y) &:= \frac{\sqrt{\pi}}{\sqrt{15}} \left(\frac{\partial^2 k(y)}{\partial x_2^2} - \frac{\partial^2 k(y)}{\partial x_1^2} \right), \\ \psi_{3,3}(y) &:= \frac{\sqrt{\pi}}{\sqrt{5}} \left(\frac{\partial^2 k(y)}{\partial x_3^2} - \frac{\Delta k(y)}{3} \right). \end{aligned}$$

Proof. To prove the claim we observe that if

$$\text{Proj}_{T_{z_{1,0}}Z^\perp} \left(\frac{1}{2} \int D^2 k(y)(x)^2 (z_{1,0})^{2^*-1} \cdot \right) = \sum_{i+j \neq 1} \sum_{l=1}^{c_i} \beta_{i,j,l} \Phi_{i,j,l}^{1,0},$$

then (2.7) implies

$$\tilde{w}_2(y) = \sum_{i+j \neq 1} \sum_{l=1}^{c_i} \frac{\beta_{i,j,l}}{\lambda_{i,j}} \Phi_{i,j,l}^{1,0}. \quad (4.14)$$

To this end we note that in the basis given in (2.6)

$$\begin{aligned} \frac{1}{2} D^2 k(y)(x)^2 &= \frac{\Delta k(y)}{6} \|x\|^2 + \sum_{l=2}^3 \psi_{l,l}(y) Y_{2,(l,l)}(x) \\ &+ \sum_{1 \leq l < m \leq 3} \psi_{l,m}(y) Y_{2,(l,m)}(x). \end{aligned} \quad (4.15)$$

Consequently,

$$\tilde{w}_2(y) \in \langle \Phi_{2,j,l}^{1,0}, \Phi_{0,j,1}^{1,0} : j \in \mathbb{N}_0, 1 \leq l \leq c_2 \rangle.$$

Using (A.1)-(A.7) we find for $n = (l, l)$ or $n = (l, m)$

$$\begin{aligned} \int \psi_n Y_{2,n}(x/|x|) |x|^2 (z_{1,0})^5 \Phi_{2,j,n}^{1,0} &= \psi_n a_{2,j} 3^{\frac{5}{4}} \frac{(j+1)\Gamma(3/2)\Gamma(j+7/2)}{2\Gamma(j+5)} \\ \int |x|^2 z_{1,0}^5 \Phi_{0,j,0}^{1,0} &= a_{0,j} 3^{\frac{5}{4}} 2\sqrt{\pi} \left(\frac{\Gamma(\frac{1}{2})\Gamma(j+\frac{3}{2})}{2\Gamma(j+2)} - \delta_{0,j} \frac{\Gamma(\frac{3}{2})^2}{2\Gamma(3)} \right). \end{aligned}$$

Now, the claim follows from (4.14) and (4.15). $\square$

Remark 4.4. Under the assumptions of Lemma 4.2 an explicit calculation together with (4.4) yields for $(t, \mu, y) \in \Omega$, where we suppress the dependence of w and $\vec{\alpha}$ on (t, μ, y) ,

$$\begin{aligned} f_t''(z_{\mu,y} + w)\varphi &= f_0''(z_{\mu,y})\varphi - 5t \int (k(x) - k(y))(z_{\mu,y} + w_0)^4 \varphi \\ &\quad - 20 \int (1 + tk(x))(z_{\mu,y} + w_0)^3 (w - w_0) \varphi \\ &\quad - 30 \int (1 + tk(x))(z_{\mu,y} + w_0)^2 (w - w_0)^2 \varphi \\ &\quad + t^3 O_{A_0}(|\nabla k(y)|^3 \min(1, \mu^3) + \min(1, \mu^6)), \end{aligned} \quad (4.16)$$

which implies by Lemma 4.1

$$\|f_t''(z_{\mu,y} + w) - f_0''(z_{\mu,y})\| \leq |t| O_{A_0}(|\nabla k(y)| \min(1, \mu) + \min(1, \mu^{\frac{7}{4}})).$$

Consequently, from (4.5), after decreasing μ_0 and t_0 if necessary,

$$\left\| \frac{\partial H}{\partial(w, \vec{\alpha})}(t, \mu, y, w, \vec{\alpha}) - \frac{\partial H}{\partial(w, \vec{\alpha})}(0, \mu, y, 0, 0) \right\| \leq \frac{1}{2} C_*$$

Thus we may assume $\frac{\partial H}{\partial(w, \vec{\alpha})}(t, \mu, y, w, \vec{\alpha})$ is invertible and its inverse is uniformly bounded with respect to $(t, \mu, y) \in \Omega$.

We begin the expansion of $\vec{\alpha}$ by computing the third order term.

Lemma 4.5. Under the assumptions of Lemma 4.2 we have as $\mu \rightarrow 0$

$$\|\vec{\alpha}(t, \mu, y) - \sum_{j=1}^3 \vec{\alpha}_j(t, \mu, y)\| = t O_{A_0}(\mu^{3+\frac{1}{2}} + |\nabla k(y)|^2 \mu^2),$$

where $\vec{\alpha}_1, \vec{\alpha}_2$ are defined in Lemma 4.2 and $\vec{\alpha}_3$ is given by

$$\vec{\alpha}_3(t, \mu, y)_i := -t\mu^3 \frac{3^{\frac{1}{4}}\pi}{2\sqrt{15}} (1 + tk(y))^{-\frac{5}{4}} \frac{\partial}{\partial x_i} \Delta k(y),$$

for $i = 1 \dots 3$ and

$$\vec{\alpha}_3(t, \mu, y)_0 := -t\mu^3 (1 + tk(y))^{-\frac{5}{4}} \frac{3^{\frac{3}{4}}4}{\pi\sqrt{5}} \oint (k(x+y) - T_{k(\cdot+y),0}^3(x)) \frac{1}{|x|^6}.$$

Proof. In the sequel we will suppress the dependence of w and $\vec{\alpha}$ on t, μ and y , when there is no possibility of confusion. Moreover, we always assume $0 < \mu \leq 1$.

As in Lemma 4.2 we infer from Lemma 4.1, (4.9), and the definition of w_1

$$\begin{aligned} f_t'(z_{\mu,y} + w) &= f_0''(z_{\mu,y})(w - w_0) - t \int (k(x) - k(y))(z_{\mu,y} + w_0)^5 \\ &\quad - 5t \int (k(x) - T_{k,y}^1(x))(z_{\mu,y} + w_0)^4 (w - w_0 - w_1) \\ &\quad - 10 \int (1 + tk(x))(z_{\mu,y} + w_0)^3 (w - w_0 - w_1)^2 \\ &\quad + t^2 O_{A_0}(|\nabla k(y)|^2 \mu^2 + |\nabla k(y)| \mu^3 + \mu^6). \end{aligned} \quad (4.17)$$

As $f'_t(z_{\mu,y} + w) - \bar{\alpha}\dot{\xi}_{\mu,y} = 0$ and $(\dot{\xi}_{\mu,y})_i \in N(f''_0(z_{\mu,y}))$ we obtain from (4.11) and testing (4.17) with $(\dot{\xi}_{\mu,y})_i$

$$\begin{aligned} (\bar{\alpha} - \sum_{j=1}^2 \bar{\alpha}_j)_i &= -t \int (k(x) - T_{k,y}^2(x)) (z_{\mu,y} + w_0)^5 (\dot{\xi}_{\mu,y})_i \\ &\quad - 5t \int (k(x) - T_{k,y}^1(x)) (z_{\mu,y} + w_0)^4 (w - w_0 - w_1) (\dot{\xi}_{\mu,y})_i \\ &\quad - 10 \int (1 + tk(x)) (z_{\mu,y} + w_0)^3 (w - w_0 - w_1)^2 (\dot{\xi}_{\mu,y})_i \\ &\quad + t^2 O_{A_0}(|\nabla k(y)|^2 \mu^2 + |\nabla k(y)| \mu^3 + \mu^6). \end{aligned} \quad (4.18)$$

By Lemma 4.1 and 4.2 we have for $0 \leq i \leq 3$

$$\begin{aligned} \int (k(x) - T_{k,y}^1(x)) (z_{\mu,y} + w_0)^4 (w - w_0 - w_1) (\dot{\xi}_{\mu,y})_i &= O_{A_0}(t\mu^{3+\frac{1}{2}}), \\ \int (1 + tk(x)) (z_{\mu,y} + w_0)^3 (w - w_0 - w_1)^2 (\dot{\xi}_{\mu,y})_i &= O_{A_0}(t\mu^4). \end{aligned} \quad (4.19)$$

If $1 \leq i \leq 3$ then we obtain from Lemma 4.1

$$\begin{aligned} \int (k(x) - T_{k,y}^2(x)) (z_{\mu,y} + w_0)^5 (\dot{\xi}_{\mu,y})_i \\ = \int \frac{1}{6} D^3 k(y) (x - y)^3 (z_{\mu,y} + w_0)^5 (\dot{\xi}_{\mu,y})_i + O_{A_0}(\mu^{3+\frac{1}{2}}), \end{aligned}$$

and from (A.9)

$$\int \frac{1}{6} D^3 k(y) (x - y)^3 (z_{\mu,y} + w_0)^5 (\dot{\xi}_{\mu,y})_i = \mu^3 \frac{3^{\frac{1}{4}} \pi}{2\sqrt{15}} (1 + tk(y))^{-\frac{5}{4}} \frac{\partial}{\partial x_i} \Delta k(y). \quad (4.20)$$

Hence, the assertion of the lemma for $1 \leq i \leq 3$ follows from (4.18)-(4.20). To treat the remaining case $i = 0$ we use the fact that $D^3 k(y)(x)^3$ is odd and get

$$\begin{aligned} \int_{B_1(y)} (k(x) - T_{k,y}^2(x)) (z_{\mu,y} + w_0)^5 (\dot{\xi}_{\mu,y})_0 \\ = \int_{B_1(y)} (k(x) - T_{k,y}^3(x)) (z_{\mu,y} + w_0)^5 (\dot{\xi}_{\mu,y})_0 \end{aligned} \quad (4.21)$$

Since, by Lemma 4.1,

$$\int_{\mathbb{R}^3} |k(x) - T_{k,y}^3(x)| \mu^{-3} (1 + \mu^{-2} |x - y|^2)^{-4} = O_{A_0}(\mu^4),$$

there holds after a translation $x \rightarrow x + y$

$$\begin{aligned} \int_{B_1(y)} (k(x) - T_{k,y}^3(x)) (z_{\mu,y} + w_0)^5 (\dot{\xi}_{\mu,y})_0 \\ = \frac{3^{\frac{3}{4}} 4 \mu^3}{\pi \sqrt{5} (1 + tk(y))^{\frac{5}{4}}} \int_{B_1(0)} \frac{(k(x + y) - T_{k(\cdot+y),0}^3(x))}{(\mu^2 + |x|^2)^3} + O_{A_0}(\mu^4). \end{aligned}$$

From

$$(\mu^2 + |x|^2)^{-3} - |x|^{-6} = -(\mu^2 + |x|^2)^{-3}(3|x|^{-2}\mu^2 + 3|x|^{-4}\mu^4 + |x|^{-6}\mu^6)$$

and Lemma 4.1, we infer

$$\int_{\mathbb{R}^3} |k(x+y) - T_{k(\cdot+y),0}^3(x)| |(\mu^2 + |x|^2)^3 - |x|^{-6}| = O_{A_0}(\mu). \quad (4.22)$$

Hence,

$$\begin{aligned} & \int (k(x) - T_{k,y}^2(x)) (z_{\mu,y} + w_0)^5 (\dot{\xi}_{\mu,y})_0 \\ &= \frac{3^{\frac{3}{4}}4\mu^3}{\pi\sqrt{5}(1+tk(y))^{\frac{5}{4}}} \left(\int_{B_1(0)} (k(x+y) - T_{k(\cdot+y),0}^3(x)) |x|^{-6} + \right. \\ & \quad \left. \int_{\mathbb{R}^3 \setminus B_1(0)} (k(x+y) - T_{k(\cdot+y),0}^2(x)) |x|^{-6} \right) + O_{A_0}(\mu^4) \\ &= \frac{3^{\frac{3}{4}}4}{\pi\sqrt{5}} \mu^3 (1+tk(y))^{-\frac{5}{4}} \oint \left(k(x+y) - T_{k(\cdot+y),0}^3(x) \right) \frac{1}{|x|^6} + O_{A_0}(\mu^4), \end{aligned}$$

which ends the proof. $\square$

Remark 4.6. From Lemma 4.2, (4.17), and (4.18) we see that

$$\frac{1}{t\mu} \vec{\alpha}(t, \mu, y)$$

is a well defined, continuous function for $(t, \mu, y) \in \Omega$.

Lemma 4.7. Under the assumptions of Lemma 4.2 we have as $\mu \rightarrow 0$

$$\begin{aligned} |\vec{\alpha}(t, \mu, y) - \sum_{j=1}^4 \vec{\alpha}_j(t, \mu, y)| &= t O_{A_0}(\mu^{4+\frac{1}{2}}) \\ &+ t^2 O_{A_0}(\mu^2 |\nabla k(y)|^2 + \mu^3 |\nabla k(y)| + \mu^4 |\Delta k(y)| + \mu^{4+\frac{1}{4}}), \end{aligned}$$

where for $1 \leq i \leq 3$

$$\vec{\alpha}_4(t, \mu, y)_i = -t\mu^4 (1+tk(y))^{-\frac{5}{4}} \frac{3^{\frac{3}{4}}8}{\pi\sqrt{5}} \oint \left(k(x+y) - T_{k(\cdot+y),0}^3(x) \right) \frac{x_i}{|x|^8}$$

and

$$\begin{aligned} \vec{\alpha}_4(t, \mu, y)_0 &= t\mu^4 (1+tk(y))^{-\frac{5}{4}} \frac{3^{\frac{3}{4}}\pi\sqrt{5}}{30} \Delta^2 k(y) \\ &- t^2 \mu^4 (1+tk(y))^{-\frac{9}{4}} \frac{3^{\frac{3}{4}}\sqrt{5}}{16} \left(\int_{\partial B_1(0)} |D^2 k(y)(x)|^2 \right). \end{aligned}$$

Proof. We proceed as in Lemma 4.5 and suppress the dependence of w and $\vec{\alpha}$ on t, μ , and y . From Lemmas 4.1 and 4.2 we have for $0 \leq i \leq 3$

$$\begin{aligned} & -5t \int (k(x) - T_{k,y}^1(x)) (z_{\mu,y} + w_0)^4 (w - w_0 - w_1) (\dot{\xi}_{\mu,y})_i \\ &= -5t \int \frac{1}{2} D^2 k(y) (x-y)^2 (z_{\mu,y} + w_0)^4 w_2 (\dot{\xi}_{\mu,y})_i + t^2 O_{A_0}(\mu^{4+\frac{1}{4}}). \end{aligned} \quad (4.23)$$

Furthermore,

$$\begin{aligned} & -10 \int (1 + tk(x))(z_{\mu,y} + w_0)^3 (w - w_0 - w_1)^2 (\dot{\xi}_{\mu,y})_i \\ & = -10 \int (1 + tk(y))(z_{\mu,y} + w_0)^3 (w_2)^2 (\dot{\xi}_{\mu,y})_i + t^2 O_{A_0}(\mu^{4+\frac{1}{4}}). \end{aligned} \quad (4.24)$$

Case $1 \leq i \leq 3$: Since w_2 is even and $(\dot{\xi}_{\mu,y})_i$ is odd for $1 \leq i \leq 3$ the integrals in (4.23)-(4.24) vanish. Thus, from (4.18) and the definition of $\vec{\alpha}_3$ we get

$$\begin{aligned} \left(\vec{\alpha} - \sum_{j=1}^3 \vec{\alpha}_j \right)_i &= -\frac{t\mu^4}{(1 + tk(y))^{\frac{5}{4}} \pi \sqrt{5}} \int \frac{(k(x+y) - T_{k(\cdot+y),0}^3(x))x_i}{(\mu^2 + |x|^2)^4} \\ & \quad + t^2 O_{A_0}(|\nabla k(y)|^2 \mu^2 + |\nabla k(y)|\mu^3 + \mu^{4+\frac{1}{4}}). \end{aligned}$$

Since $D^4 k(y)(x)^4$ is even we may proceed analogously as in (4.21) and prove the claim of the lemma if $1 \leq i \leq 3$.

Case $i = 0$: From (4.18), the definition of $\vec{\alpha}_3$, (4.22), (4.23)-(4.24), and the fact that $D^3 k(y)(x)^3$ is odd, we arrive at

$$\begin{aligned} & (\vec{\alpha} - \vec{\alpha}_1 - \vec{\alpha}_2 - \vec{\alpha}_3)_0 \\ &= \frac{3^{\frac{3}{4}} 4t}{\pi \sqrt{5} (1 + tk(y))^{\frac{5}{4}}} \int (k(x+y) - T_{k(\cdot+y),0}^3(x)) \\ & \quad \left(\frac{3|\frac{x}{\mu}|^{-2} + 3|\frac{x}{\mu}|^{-4} + |\frac{x}{\mu}|^{-6}}{\mu^3(1 + |\frac{x}{\mu}|^2)^3} + \frac{2}{\mu^3(1 + |\frac{x}{\mu}|^2)^4} \right) \\ & \quad - 5t \int \frac{1}{2} D^2 k(y)(x-y)^2 (z_{\mu,y} + w_0)^4 w_2 (\dot{\xi}_{\mu,y})_0 \\ & \quad - 10 \int (1 + tk(y))(z_{\mu,y} + w_0)^3 (w_2)^2 (\dot{\xi}_{\mu,y})_0 \\ & \quad + t^2 O_{A_0}(|\nabla k(y)|^2 \mu^2 + |\nabla k(y)|\mu^3 + \mu^{4+\frac{1}{4}}). \end{aligned} \quad (4.25)$$

From Lemma 4.1, (A.6), and (A.10) we get

$$\begin{aligned} & \int (k(x+y) - T_{k(\cdot+y),0}^3(x)) \left(\frac{3|\frac{x}{\mu}|^{-2} + 3|\frac{x}{\mu}|^{-4} + |\frac{x}{\mu}|^{-6}}{\mu^3(1 + |\frac{x}{\mu}|^2)^3} + \frac{2}{\mu^3(1 + |\frac{x}{\mu}|^2)^4} \right) \\ &= \mu^4 \int \frac{1}{4!} D^4 k(y)(x)^4 \frac{5 + 6|x|^{-2} + 4|x|^{-4} + |x|^{-6}}{(1 + |x|^2)^4} + O_{A_0}(\mu^{4+\frac{1}{2}}) \\ &= \mu^4 \frac{\pi^2}{24} \Delta^2 k(y) + O_{A_0}(\mu^{4+\frac{1}{2}}). \end{aligned}$$

For the second term in (4.25) we obtain from Lemmas 4.2 and 4.3

$$\begin{aligned} & -5t \int \frac{1}{2} D^2 k(y)(x-y)^2 (z_{\mu,y} + w_0)^4 w_2 (\dot{\xi}_{\mu,y})_0 \\ &= -\frac{t^2 \mu^4 4\sqrt{15}}{\pi(1 + tk(y))^{\frac{9}{4}}} \int \frac{1}{2} D^2 k(y)(x)^2 \tilde{w}_2(y) (1 + |x|^2)^{-\frac{5}{2}} \left(1 - \frac{2}{1 + |x|^2} \right). \end{aligned}$$

Moreover, from Lemma 4.3, (4.15), and (A.12)

$$\begin{aligned}
& \int \frac{1}{2} D^2 k(y) (x)^2 \tilde{w}_2(y) (1 + |x|^2)^{-\frac{5}{2}} (1 - 2(1 + |x|^2)^{-1}) \\
&= O_{A_0}(|\Delta k(y)|^2) + \left(\sum_{1 \leq l < m \leq 3} \psi_{l,m}(y)^2 + \sum_{l=2}^3 \psi_{l,l}(y)^2 \right) \frac{3^{\frac{5}{4}} \sqrt{\pi}}{4} \\
&\quad \times \sum_{j=0}^{\infty} \frac{(3+j)j!}{\Gamma(\frac{7}{2}+j)} \int_0^{\infty} \frac{r^6}{(1+r^2)^5} \left(1 - \frac{2}{1+r^2}\right) \mathcal{P}_j^{(\frac{5}{2}, \frac{5}{2})} \left(1 - \frac{2}{1+r^2}\right) \\
&= O_{A_0}(|\Delta k(y)|^2) + \frac{3^{\frac{5}{4}} \pi}{64} \left(\int_{\partial B_1(0)} |D^2 k(y)(x)^2 - \frac{\Delta k(y)}{3} |x|^2|^2 \right) \\
&\quad \times \sum_{j=0}^{\infty} \frac{2+6j+j^2}{(5+j)(4+j)(2+j)(1+j)} \\
&= O_{A_0}(|\Delta k(y)|) + \frac{3^{\frac{5}{4}} 5 \pi}{1024} \left(\int_{\partial B_1(0)} |D^2 k(y)(x)^2|^2 \right). \tag{4.26}
\end{aligned}$$

For the remaining term in (4.25) we derive

$$\begin{aligned}
& -10 \int (1 + tk(y))(z_{\mu,y} + w_0)^3 (w_2)^2 (\dot{\xi}_{\mu,y})_0 \\
&= -10 \frac{t^2 \mu^4}{(1 + tk(y))^{\frac{9}{4}}} \frac{3^{\frac{3}{4}} 4}{\pi \sqrt{15}} \int (1 + |x|^2)^{-2} \left(1 - \frac{2}{1 + |x|^2}\right) (\tilde{w}_2(y))^2.
\end{aligned}$$

By Lemma 4.3, (4.26), and we have with $\beta_j := \frac{(3+j)j!}{\Gamma(j+\frac{7}{2})}$

$$\begin{aligned}
& \int (1 + |x|^2)^{-2} \left(1 - \frac{2}{1 + |x|^2}\right) (\tilde{w}_2(y))^2 \\
&= O_{A_0}(|\Delta k(y)|) + \frac{1}{4} \left(\int_{\partial B_1(0)} |D^2 k(y)(x)^2|^2 \right) \frac{3^{\frac{5}{2}} \pi}{16} \\
&\quad \times \int_0^{\infty} r^6 (1 + r^2)^{-7} \left(1 - \frac{2}{1 + r^2}\right) \left(\sum_{j=0}^{\infty} \beta_j \mathcal{P}_j^{(\frac{5}{2}, \frac{5}{2})} \left(1 - \frac{2}{1 + r^2}\right) \right)^2 \\
&= \frac{1}{4} \left(\int_{\partial B_1(0)} |D^2 k(y)(x)^2|^2 \right) \frac{3^{\frac{5}{2}} \pi}{16} \frac{1}{288} + O_{A_0}(|\Delta k(y)|). \tag{4.27}
\end{aligned}$$

Combining the computations in (4.25)- (4.27) ends the proof. $\square$

Lemma 4.8. *Under the assumptions of Lemma 4.2 suppose $k \in C^6(\mathbb{R}^3)$ and $\|D^6 k\|_{\infty} \leq A_0$. Then there holds*

$$\begin{aligned}
& \|(\vec{\alpha}(t, \mu, y))_0 - \sum_{j=1}^5 (\vec{\alpha}_j(t, \mu, y))_0\| = O_{A_0}(t\mu^6) \\
& \quad + t^2 O_{A_0} \left(\mu^2 |\nabla k(y)|^2 + \mu^3 |\nabla k(y)| + \mu^4 |\Delta k(y)| + \mu^{4+\frac{1}{4}} \right).
\end{aligned}$$

where $(\vec{\alpha}_5(t, \mu, y))_0$ is given by

$$(\vec{\alpha}_5(t, \mu, y))_0 := -\frac{t\mu^5}{(1+tk(y))^{\frac{5}{4}}} \frac{3^{\frac{3}{4}}4\sqrt{5}}{\pi} \oint \left(k(y+x) - T_{k(\cdot+y),0}^4(x) \right) \frac{1}{|x|^8}.$$

Proof. From the proof of Lemma 4.7 we infer

$$\begin{aligned} (\vec{\alpha}(t, \mu, y))_0 - \sum_{j=1}^4 (\vec{\alpha}_j(t, \mu, y))_0 &= -t(1+tk(y))^{-\frac{5}{4}} \frac{3^{\frac{5}{4}}4}{\pi\sqrt{15}} \\ &\int \left(k(y+x) - \sum_{\ell=0}^4 \frac{1}{\ell!} D^\ell k(y)(x)^\ell \right) \frac{5+6|\frac{x}{\mu}|^{-2}+4|\frac{x}{\mu}|^{-4}+|\frac{x}{\mu}|^{-6}}{\mu^3 \left(1+|\frac{x}{\mu}|^2\right)^4} \\ &+ t^2 O_{A_0}(\mu^2 |\nabla k(y)|^2 + \mu^3 |\nabla k(y)| + \mu^4 |\Delta k(y)| + \mu^{4+\frac{1}{4}}) \end{aligned}$$

Since $D^5 k(y)(x)^5$ is odd, analogously as in Lemma 4.5 we find

$$\begin{aligned} &\int \left(k(y+x) - \sum_{\ell=0}^4 \frac{1}{\ell!} D^\ell k(y)(x)^\ell \right) \frac{5+6|\frac{x}{\mu}|^{-2}+4|\frac{x}{\mu}|^{-4}+|\frac{x}{\mu}|^{-6}}{\mu^3 \left(1+|\frac{x}{\mu}|^2\right)^4} \\ &= 5\mu^5 \oint \left(k(y+x) - \sum_{\ell=0}^4 \frac{1}{\ell!} D^\ell k(y)(x)^\ell \right) \frac{1}{|x|^8} + O_{A_0}(\mu^6), \end{aligned}$$

which ends the proof. $\square$

5. DERIVATIVES OF $\vec{\alpha}$

Lemma 5.1. *Under the assumptions of Lemma 4.2 we have for all $(t, \mu, y) \in \Omega$ with $|\mu| \leq 1$ and $1 \leq i, j \leq 3$*

$$\left| \frac{1}{t\mu} \frac{\partial \alpha(t, \mu, y)_i}{\partial y_j} + \frac{\pi}{3^{\frac{1}{4}}\sqrt{5}} (1+tk(y))^{-\frac{5}{4}} \frac{\partial^2 k(y)}{\partial x_i \partial x_j} \right| \leq O_{A_0}(|\nabla k(y)|^2 + \mu^{1+\frac{1}{4}}), \quad (5.1)$$

$$\begin{aligned} &\left| \frac{1}{t\mu^2} \frac{\partial \alpha(t, \mu, y)_0}{\partial y_j} + \frac{\pi}{3^{\frac{1}{4}}\sqrt{5}} (1+tk(y))^{-\frac{5}{4}} \frac{\partial}{\partial x_j} \Delta k(y) \right| \\ &\leq O_{A_0}(|\nabla k(y)|^2 \mu^{-1} + \mu^{\frac{1}{4}}). \end{aligned} \quad (5.2)$$

Proof. In the sequel we will suppress the dependence of w and $\vec{\alpha}$ on (t, μ, y) . Since $H(t, \mu, y, w, \vec{\alpha}) \equiv 0$ we have

$$-\frac{\partial H}{\partial y} = \frac{\partial H}{\partial(w, \vec{\alpha})} \begin{pmatrix} \frac{\partial w}{\partial y} \\ \frac{\partial \vec{\alpha}}{\partial y} \end{pmatrix} = \frac{\partial H}{\partial w} \frac{\partial w}{\partial y} + \frac{\partial H}{\partial \vec{\alpha}} \frac{\partial \vec{\alpha}}{\partial y}, \quad (5.3)$$

where

$$\frac{\partial H}{\partial y_j}(t, \mu, y, w, \vec{\alpha}) = \left(f_t''(z_{\mu,y} + w) \frac{\partial z_{\mu,y}}{\partial y_j} - \vec{\alpha} \cdot \frac{\partial \dot{\xi}_{\mu,y}}{\partial y_j}, \left(\langle w, \frac{\partial(\dot{\xi}_{\mu,y})_l}{\partial y_j} \rangle \right)_l \right). \quad (5.4)$$

A direct calculation gives for $0 \leq l \leq 3$

$$\left\| \frac{\partial(\dot{\xi}_{\mu,y})_l}{\partial y_j} \right\| + \left\| \frac{\partial z_{\mu,y}}{\partial y_j} \right\| \leq \text{const } \mu^{-1}. \quad (5.5)$$

Differentiating $\langle z_{\mu,y}, (\dot{\xi}_{\mu,y})_l \rangle \equiv 0$ leads to

$$\langle z_{\mu,y}, \frac{\partial(\dot{\xi}_{\mu,y})_l}{\partial y_j} \rangle = \langle \frac{\partial z_{\mu,y}}{\partial y_j}, (\dot{\xi}_{\mu,y})_l \rangle,$$

and with (4.4) and (5.5) we arrive at

$$\left| \langle w, \frac{\partial(\dot{\xi}_{\mu,y})_l}{\partial y_j} \rangle - ((1 + tk(y))^{-\frac{1}{4}} - 1) \langle \frac{\partial z_{\mu,y}}{\partial y_j}, (\dot{\xi}_{\mu,y})_l \rangle \right| = tO_{A_0}(|\nabla k(y)| + \mu). \quad (5.6)$$

By (5.5), the expansion of $\vec{\alpha}$ in Lemma 4.2, and (4.16) we see

$$\left\| f_t''(z_{\mu,y} + w) \frac{\partial z_{\mu,y}}{\partial y_j} \right\| + \left\| \vec{\alpha} \cdot \frac{\partial \dot{\xi}_{\mu,y}}{\partial y_j} \right\| \leq tO_{A_0}(|\nabla k(y)| + \mu). \quad (5.7)$$

From (5.6)-(5.7) we get

$$\left\| \frac{\partial H}{\partial(w, \vec{\alpha})}(t, \mu, y, w, \vec{\alpha}) \begin{pmatrix} ((1 + tk(y))^{-\frac{1}{4}} - 1) \frac{\partial z_{\mu,y}}{\partial y_j} \\ 0 \end{pmatrix} + \frac{\partial H}{\partial y_j}(t, \mu, y, w, \vec{\alpha}) \right\| \leq tO_{A_0}(|\nabla k(y)| + \mu)$$

which implies due to the uniform bound of the inverse (see Remark 4.4)

$$\left\| \frac{\partial w}{\partial y_j} - ((1 + tk(y))^{-\frac{1}{4}} - 1) \frac{\partial z_{\mu,y}}{\partial y_j} \right\| + \left\| \frac{\partial \vec{\alpha}}{\partial y_j} \right\| \leq tO_{A_0}(|\nabla k(y)| + \mu). \quad (5.8)$$

From (4.5) and (5.3)-(5.4) we deduce after testing with $(\dot{\xi}_{\mu,y})_j$

$$\frac{\partial \vec{\alpha}_j}{\partial y_i} = f_t''(z_{\mu,y} + w) \left(\frac{\partial z_{\mu,y}}{\partial y_i} + \frac{\partial w}{\partial y_i} \right) (\dot{\xi}_{\mu,y})_j - \sum_{l=0}^3 \vec{\alpha}(t, \mu, y)_l \langle \frac{\partial(\dot{\xi}_{\mu,y})_l}{\partial y_i}, (\dot{\xi}_{\mu,y})_j \rangle. \quad (5.9)$$

From Lemma 4.1, (4.16), (5.8), and the fact that $(\dot{\xi}_{\mu,y})_j \in N(f_0''(z_{\mu,y}))$ we obtain

$$\begin{aligned} & f_t''(z_{\mu,y} + w) \left(\frac{\partial z_{\mu,y}}{\partial y_i} + \frac{\partial w}{\partial y_i} \right) (\dot{\xi}_{\mu,y})_j \\ &= -\frac{5t}{(1 + tk(y))^{\frac{5}{4}}} \int \left(\sum_{\ell=1}^3 \frac{1}{\ell!} D^\ell k(y) (x - y)^\ell \right) (z_{\mu,y})^4 \left(\frac{\partial z_{\mu,y}}{\partial y_i} \right) (\dot{\xi}_{\mu,y})_j \\ &\quad - 20 \int (z_{\mu,y})^3 (w - w_0) \left(\frac{\partial z_{\mu,y}}{\partial y_i} \right) (\dot{\xi}_{\mu,y})_j \\ &\quad + O_{A_0} \left(t\mu^{2+\frac{3}{4}} + t^2(|\nabla k(y)|^2 \mu + |\nabla k(y)| \mu^{\frac{7}{4}} + \mu^{2+\frac{3}{4}}) \right) \end{aligned} \quad (5.10)$$

Differentiating the identity $f_0''(z_{\mu,y})(\dot{\xi}_{\mu,y})_j = 0$ with respect to y_i leads to

$$0 = f_0'''(z_{\mu,y}) \left(\frac{\partial z_{\mu,y}}{\partial y_i} \right) (\dot{\xi}_{\mu,y})_j + f_0''(z_{\mu,y}) \left(\frac{\partial(\dot{\xi}_{\mu,y})_j}{\partial y_i} \right),$$

and we get from Lemma 4.1, (4.2), and (4.17)

$$\begin{aligned}
& -20 \int (z_{\mu,y})^3 (w - w_0) \frac{\partial z_{\mu,y}}{\partial y_i} (\dot{\xi}_{\mu,y})_j \\
& = f_0'''(z_{\mu,y})(w - w_0) \frac{\partial z_{\mu,y}}{\partial y_i} (\dot{\xi}_{\mu,y})_j = -f_0''(z_{\mu,y}) \frac{\partial (\dot{\xi}_{\mu,y})_j}{\partial y_i} (w - w_0) \\
& = - \sum_{l=0}^3 (\vec{\alpha}(t, \mu, y))_l \langle (\dot{\xi}_{\mu,y})_l, \frac{\partial (\dot{\xi}_{\mu,y})_j}{\partial y_i} \rangle \\
& \quad - \frac{t}{(1 + tk(y))^{\frac{5}{4}}} \int \sum_{\ell=1}^3 \frac{1}{\ell!} D^\ell k(y) (x - y)^\ell (z_{\mu,y})^5 \frac{\partial (\dot{\xi}_{\mu,y})_j}{\partial y_i} \\
& \quad + tO_{A_0}(\mu |\nabla k(y)|^2 + \mu^{2+\frac{1}{4}}).
\end{aligned}$$

Differentiating $\langle (\dot{\xi}_{\mu,y})_l, (\dot{\xi}_{\mu,y})_j \rangle \equiv \text{const}$ with respect to y_i we obtain

$$\left\langle \frac{\partial (\dot{\xi}_{\mu,y})_l}{\partial y_i}, (\dot{\xi}_{\mu,y})_j \right\rangle = - \left\langle (\dot{\xi}_{\mu,y})_l, \frac{\partial (\dot{\xi}_{\mu,y})_j}{\partial y_i} \right\rangle.$$

Inserting the above computations in (5.10) and (5.9) leads to

$$\begin{aligned}
\frac{\partial \vec{\alpha}_j}{\partial y_i} & = - \frac{t}{(1 + tk(y))^{\frac{5}{4}}} \int \left(\sum_{\ell=1}^3 \frac{1}{\ell!} D^\ell k(y) (x - y)^\ell \right) \frac{d}{dy_i} \left((z_{\mu,y})^5 (\dot{\xi}_{\mu,y})_j \right) \\
& \quad + tO_{A_0}(|\nabla k(y)|^2 \mu + \mu^{2+\frac{1}{4}}).
\end{aligned}$$

As $\frac{\partial}{\partial y_i} z_{\mu,y}(x) = -\frac{\partial}{\partial x_i} z_{\mu,y}(x)$ and $\frac{\partial}{\partial y_i} (\dot{\xi}_{\mu,y})_j(x) = -\frac{\partial}{\partial x_i} (\dot{\xi}_{\mu,y})_j(x)$ we obtain by partial integration

$$\begin{aligned}
& \int \left(\sum_{\ell=1}^3 \frac{1}{\ell!} D^\ell k(y) (x - y)^\ell \right) \frac{d}{dy_i} \left((z_{\mu,y})^5 (\dot{\xi}_{\mu,y})_j \right) \\
& = \int \frac{d}{dx_i} \left(\sum_{\ell=1}^3 \frac{1}{\ell!} D^\ell k(y) (x - y)^\ell \right) (z_{\mu,y})^5 (\dot{\xi}_{\mu,y})_j \\
& = \int \left(\sum_{\ell=0}^2 \frac{1}{\ell!} D^\ell \frac{\partial k}{\partial x_i}(y) (x - y)^\ell \right) (z_{\mu,y})^5 (\dot{\xi}_{\mu,y})_j.
\end{aligned}$$

The latter integral may be evaluated as in Lemma 4.2 and yields the claim. $\square$

Lemma 5.2. *Under the assumptions of Lemma 5.1 we have*

$$\left\| \frac{\partial w}{\partial t} - \sum_{i=0}^2 \frac{\partial w_i}{\partial t} \right\| + \left\| \frac{\partial \vec{\alpha}}{\partial t} - \sum_{i=1}^2 \frac{\partial \vec{\alpha}_i}{\partial t} \right\| = O_{A_0}(|\nabla k(y)|^2 \mu^2 + \mu^{2+\frac{1}{4}}). \quad (5.11)$$

Proof. In the sequel we will suppress the dependence of w and $\vec{\alpha}$ on (t, μ, y) . Since $H(t, \mu, y, w, \vec{\alpha}) \equiv 0$ we have

$$\begin{aligned}
\left(\int k(x) (z_{\mu,y} + w)^5 \cdot, \vec{0} \right) & = - \frac{\partial H}{\partial t} \Big|_{(t, \mu, y, w, \vec{\alpha})} = \frac{\partial H}{\partial (w, \vec{\alpha})} \Big|_{(t, \mu, y, w, \vec{\alpha})} \begin{pmatrix} \frac{\partial w}{\partial t} \\ \frac{\partial \vec{\alpha}}{\partial t} \end{pmatrix} \\
& = \left(f_t''(z_{\mu,y} + w) \frac{\partial w}{\partial t} - \frac{\partial \vec{\alpha}}{\partial t} \cdot \dot{\xi}_{\mu,y}, \left(\left\langle \frac{\partial w}{\partial t}, (\dot{\xi}_{\mu,y})_l \right\rangle \right)_l \right). \quad (5.12)
\end{aligned}$$

Differentiating the identities

$$f'_0(z_{\mu,y} + w_0) = t \int k(y)(z_{\mu,y} + w_0)^5, \\ f''_0(z_{\mu,y}) \sum_{i=1}^2 w_i - \sum_{i=1}^2 \vec{\alpha}_i \cdot \dot{\xi}_{\mu,y} = t \int \left(\sum_{\ell=1}^2 \frac{1}{\ell!} D^\ell k(y)(x-y)^\ell \right) (z_{\mu,y} + w_0)^5 \cdot$$

with respect to t leads to

$$f''_0(z_{\mu,y}) \sum_{i=0}^2 \frac{\partial w_i}{\partial t} - \sum_{i=1}^2 \frac{\partial \vec{\alpha}_i}{\partial t} \cdot \dot{\xi}_{\mu,y} = \int \left(\sum_{\ell=0}^2 \frac{1}{\ell!} D^\ell k(y)(x-y)^\ell \right) (z_{\mu,y} + w_0)^5 \cdot \\ + 5t \int \left(\sum_{\ell=1}^2 \frac{1}{\ell!} D^\ell k(y)(x-y)^\ell \right) (z_{\mu,y} + w_0)^4 \frac{\partial w_0}{\partial t} \cdot \quad (5.13)$$

Furthermore, we note that

$$\frac{\partial w_0}{\partial t} = -\frac{k(y)}{4} (1 + tk(y))^{-1} (z_{\mu,y} + w_0). \quad (5.14)$$

For $\frac{\partial w_i}{\partial t} = O_{A_0}(\mu^i)$ as $\mu \rightarrow 0$ we get from Lemma 4.1, (4.16), (5.13), and (5.14)

$$f''_t(z_{\mu,y} + w) \sum_{i=0}^2 \frac{\partial w_i}{\partial t} - \frac{\partial(\vec{\alpha}_1 + \vec{\alpha}_2)}{\partial t} \cdot \dot{\xi}_{\mu,y} \\ = \int \left(\sum_{\ell=0}^2 \frac{1}{\ell!} D^\ell k(y)(x-y)^\ell \right) (z_{\mu,y} + w_0)^5 \cdot \\ + 5 \int k(y)(z_{\mu,y} + w_0)^4 (w - w_0) \cdot + t O_{A_0}(|\nabla k(y)|^2 \mu^2 + \mu^{\frac{9}{4}}). \quad (5.15)$$

Moreover, by Lemmas 4.1 and 4.2

$$\int k(x)(z_{\mu,y} + w)^5 \cdot \\ = \int \left(\sum_{\ell=0}^2 \frac{1}{\ell!} D^\ell k(y)(x-y)^\ell \right) (z_{\mu,y} + w_0)^5 \cdot + O_{A_0}(\mu^{\frac{9}{4}}) \\ + 5 \int k(y)(z_{\mu,y} + w_0)^4 (w - w_0) \cdot + t O_{A_0}(|\nabla k(y)|^2 \mu^2 + \mu^3). \quad (5.16)$$

Combining (5.15), (5.16) and the fact that $\frac{\partial w_i}{\partial t}$ remains in $T_{z_{\mu,y}} Z^\perp$ we get

$$\frac{\partial H}{\partial(w, \vec{\alpha})} \left(\sum_{i=0}^2 \frac{\partial w_i}{\partial t} \right) = \frac{\partial H}{\partial t} + O_{A_0}(|\nabla k(y)|^2 \mu^2 + \mu^{2+\frac{1}{4}}).$$

and the claim of the lemma follows from Remark 4.4. $\square$

Lemma 5.3. *Under the assumptions of Lemma 5.1 we have*

$$\frac{\partial(\vec{\alpha})_0}{\partial t} = \frac{1}{t}(\vec{\alpha})_0 + \sum_{j=2}^4 \frac{\partial(\vec{\alpha}_j)_0}{\partial t} - \frac{1}{t}(\vec{\alpha}_j)_0 \\ + t O_{A_0}(|\nabla k(y)|^2 \mu^2 + |\nabla k(y)| \mu^3 + \mu^4 |\Delta k(y)|^2 + \mu^{4+\frac{1}{4}}), \quad (5.17)$$

and for $1 \leq i \leq 3$

$$\begin{aligned} \frac{\partial(\vec{\alpha})_i}{\partial t} &= \frac{1}{t}(\vec{\alpha})_i + \sum_{j=1}^3 \frac{\partial(\vec{\alpha}_j)_i}{\partial t} - \frac{1}{t}(\vec{\alpha}_j)_i \\ &\quad + tO_{A_0}(|\nabla k(y)|^2\mu^2 + |\nabla k(y)|\mu^3 + \mu^{3+\frac{1}{2}}). \end{aligned} \quad (5.18)$$

Proof. By (5.12) we have for $0 \leq i \leq 3$

$$\frac{\partial(\vec{\alpha})_i}{\partial t} = f_t''(z_{\mu,y} + w) \frac{\partial w}{\partial t} (\dot{\xi}_{\mu,y})_i - \int k(x)(z_{\mu,y} + w)^5 (\dot{\xi}_{\mu,y})_i.$$

To prove the claim of the lemma we will proceed termwise. In the calculations below certain terms will vanish simply because we are integrating a product of an odd and an even function. Moreover, we often use Lemma 4.1 without mentioning it explicitly.

For $(\dot{\xi}_{\mu,y})_i \in N(f_0''(z_{\mu,y}))$ and by (4.16) we see

$$\begin{aligned} f_t''(z_{\mu,y} + w) \frac{\partial w}{\partial t} (\dot{\xi}_{\mu,y})_i &= -5t \int (k(x) - k(y))(z_{\mu,y} + w_0)^4 \frac{\partial w}{\partial t} (\dot{\xi}_{\mu,y})_i \\ &\quad - 20 \int (1 + tk(x))(z_{\mu,y} + w_0)^3 (w - w_0) \frac{\partial w}{\partial t} (\dot{\xi}_{\mu,y})_i \\ &\quad - 30 \int (1 + tk(x))(z_{\mu,y} + w_0)^2 (w - w_0)^2 \frac{\partial w}{\partial t} (\dot{\xi}_{\mu,y})_i \\ &\quad + t^3 O_{A_0}(|\nabla k(y)|^3 \mu^3 + \mu^6). \end{aligned}$$

Due to Lemma 5.2 we may replace $\frac{\partial w}{\partial t}$ by $\sum_{i=0}^2 \frac{\partial w_i}{\partial t}$. By (5.14) we obtain

$$\begin{aligned} &-5t \int (k(x) - k(y))(z_{\mu,y} + w_0)^4 \frac{\partial w}{\partial t} (\dot{\xi}_{\mu,y})_i \\ &= \frac{5k(y)}{4(1 + tk(y))} t \int (k(x) - k(y))(z_{\mu,y} + w_0)^5 (\dot{\xi}_{\mu,y})_i \\ &\quad - 5t \int \frac{1}{2} D^2 k(y) (x - y)^2 (z_{\mu,y} + w_0)^4 \frac{\partial w_2}{\partial t} (\dot{\xi}_{\mu,y})_i \\ &\quad + tO_{A_0}(\mu^2 |\nabla k(y)|^2 + \mu^3 |\nabla k(y)| + \mu^{4+\frac{1}{4}}). \end{aligned}$$

From (5.14) and as $(\dot{\xi}_{\mu,y})_i \in N(f_0''(z_{\mu,y}))$ and $w - w_0 \in T_{z_{\mu,y}} T^\perp$ we see

$$\begin{aligned} &-20 \int (1 + tk(y))(z_{\mu,y} + w_0)^3 (w - w_0) \frac{\partial w_0}{\partial t} (\dot{\xi}_{\mu,y})_i \\ &= \frac{k(y) \langle (w - w_0), (\dot{\xi}_{\mu,y})_i \rangle}{1 + tk(y)} = 0. \end{aligned}$$

By Lemma 4.1 and (5.11) we now get

$$\begin{aligned} &-20 \int (1 + tk(x))(z_{\mu,y} + w_0)^3 (w - w_0) \frac{\partial w}{\partial t} (\dot{\xi}_{\mu,y})_i \\ &= -20 \int (1 + tk(y))(z_{\mu,y} + w_0)^3 w_2 \frac{\partial w_2}{\partial t} (\dot{\xi}_{\mu,y})_0 \\ &\quad - 20t \int \frac{1}{2} D^2 k(y) (x - y)^2 (z_{\mu,y} + w_0)^3 w_2 \frac{\partial w_0}{\partial t} (\dot{\xi}_{\mu,y})_i \\ &\quad + tO_{A_0}(\mu^2 |\nabla k(y)|^2 + \mu^3 |\nabla k(y)| + \mu^{4+\frac{1}{4}}). \end{aligned}$$

Furthermore, we see

$$\begin{aligned} & -30 \int (1 + tk(x))(z_{\mu,y} + w_0)^2 (w - w_0)^2 \frac{\partial w}{\partial t} (\dot{\xi}_{\mu,y})_i \\ & = -30 \int (1 + tk(y))(z_{\mu,y} + w_0)^2 (w_2)^2 \frac{\partial w_0}{\partial t} (\dot{\xi}_{\mu,y})_i \\ & \quad + t^2 O_{A_0}(\mu^2 |\nabla k(y)|^2 + \mu^3 |\nabla k(y)| + \mu^{4+\frac{1}{4}}) \end{aligned}$$

Since $z_{\mu,y}$ and w are orthogonal to $(\dot{\xi}_{\mu,y})_i$ and $(\dot{\xi}_{\mu,y})_i \in N(f_0''(z_{\mu,y}))$, we may estimate using Lemma 4.2

$$\begin{aligned} - \int k(x)(z_{\mu,y} + w)^5 (\dot{\xi}_{\mu,y})_i &= \frac{1}{t} f_t'(z_{\mu,y} + w) (\dot{\xi}_{\mu,y})_i + \frac{1}{t} \int (z_{\mu,y} + w)^5 (\dot{\xi}_{\mu,y})_i \\ &= \frac{1}{t} (\vec{\alpha})_i + \frac{10}{t} \int (z_{\mu,y} + w_0)^3 (w_2)^2 (\dot{\xi}_{\mu,y})_i \\ &\quad + t^2 O_{A_0}(|\nabla k(y)|^2 \mu^2 + |\nabla k(y)| \mu^3 + \mu^{4+\frac{1}{4}}). \end{aligned}$$

As

$$\frac{\partial w_2}{\partial t} = \left(\frac{1}{t} - \frac{5k(y)}{4(1 + tk(y))} \right) w_2,$$

we end up with integrals that are, up to a factor, computed in Section 4. Summing up the results will give the claim of the lemma. $\square$

6. SOLVABILITY OF $\vec{\alpha}(t, \mu, y) = 0$

Lemma 6.1. *Under the assumptions of Lemma 4.2 suppose y_0 is a nondegenerate critical point of k , i.e.*

$$\nabla k(y_0) = 0 \text{ and } D^2 k(y_0) \text{ is invertible, with } \|(D^2 k(y_0))^{-1}\| \leq A_0.$$

Moreover, assume $\Delta k(y_0) = 0$. Consider the function $\hat{\alpha}$, defined by

$$\hat{\alpha}(t, \mu, y) := \frac{3^{\frac{1}{4}} \sqrt{5}}{t\mu\pi} (1 + tk(y_0))^{\frac{5}{4}} (\vec{\alpha}(t, \mu, y)_1, \dots, \vec{\alpha}(t, \mu, y)_3)^T,$$

which is well defined and continuous in Ω (see Remark 4.6), analogously we define $\hat{\alpha}_j(t, \mu, y)$. Then there are $\delta_1 = \delta_1(A_0) > 0$ and a C^2 -function β ,

$$\beta : \{(t, \mu) : t \in [-B_0, B_1], 0 < \mu < \delta_1\} \rightarrow \mathbb{R}^3,$$

such that

$$\hat{\alpha}(t, \mu, \beta(t, \mu)) = 0 \text{ for all } t \in [-B_0, B_1], 0 < \mu < \delta_1,$$

and

$$\beta(t, \mu) = y_0 + (D^2 k(y_0))^{-1} \left(\sum_{j=3}^4 \hat{\alpha}_j(t, \mu, y_0) \right) + O_{A_0}(\mu^{3+\frac{1}{4}}).$$

Moreover, β is unique in the sense that, if $y \in B_{\delta_1}(y_0)$ satisfies $\hat{\alpha}(t, \mu, y) = 0$ for some $t \in [-B_0, B_1]$ and $0 < \mu < \delta_1$, then $y = \beta(t, \mu)$.

Proof. In view of (5.1) we would like to apply the implicit function theorem to the function $(\vec{\alpha}(t, \mu, y))_{1 \leq i \leq 3}$ in the point $(t, 0, y_0)$, but unfortunately $\vec{\alpha}$ may not be differentiable for $\mu = 0$. Instead we mimic the proof of the implicit function theorem and apply Banach's fixed-point theorem to the function

$$F(t, \mu, y) := y + (D^2k(y_0))^{-1} \hat{\alpha}(t, \mu, y)$$

in $B_\delta(y_0)$, where $\delta > 0$ will be chosen later. Fix $y_1, y_2 \in B_\delta(y_0)$, then by Lemma 5.1

$$\begin{aligned} & |F(t, \mu, y_1) - F(t, \mu, y_2)| \\ &= |(y_1 - y_2) + (D^2k(y_0))^{-1} \int_0^1 \frac{\partial \hat{\alpha}}{\partial y}(t, \mu, y_2 + t(y_1 - y_2))(y_1 - y_2) dt| \\ &\leq |(y_1 - y_2) - \left(\int_0^1 (D^2k(y_0))^{-1} D^2k(y_2 + t(y_1 - y_2)) dt \right) (y_1 - y_2)| \\ &\quad + O_{A_0} \left(\sup_{y \in B_\delta(y_0)} |\nabla k(y)| + \mu^{\frac{1}{4}} \right) |y_1 - y_2| \\ &\leq O_{A_0} \left(\delta + \mu^{\frac{1}{4}} \right) |y_1 - y_2|. \end{aligned}$$

For $y \in B_\delta(y_0)$ we estimate using Lemma 4.2

$$\begin{aligned} |F(t, \mu, y) - y_0| &= |y - y_0 + (D^2k(y_0))^{-1} (\hat{\alpha}(t, \mu, y))| \\ &\leq \left| y - y_0 - (D^2k(y_0))^{-1} (\nabla k(y) + O_{A_0}(\mu^2)) \right| \\ &\leq O_{A_0}(\delta^2 + \mu^2). \end{aligned}$$

Consequently, there is $\delta_1 = \delta_1(A_0) > 0$ such that $F(t, \mu, \cdot)$ is a contraction in $B_{\delta_1}(y_0)$ for any $0 < \mu < \delta_1$ and $t \in [-B_0, B_1]$. From Banach's fixed-point theorem we may define $\beta(t, \mu)$ to be the unique fixed-point of $F(t, \mu, \cdot)$ in $B_{\delta_1}(y_0)$. After shrinking δ_1 if necessary we may apply Lemma 5.1 and the usual implicit function theorem to see that the function β is twice differentiable for $\mu > 0$.

To deduce the expansion for small μ we fix $\rho > 0$ and

$$y \in U_\rho := B_\rho \left(y_0 + (D^2k(y_0))^{-1} \left(\sum_{j=3}^4 \hat{\alpha}_j(t, \mu, y_0) \right) \right).$$

Then, by Lemmas 4.5 and 4.7

$$\begin{aligned} & \left| F(t, \mu, y) - y_0 - (D^2k(y_0))^{-1} \left(\sum_{j=3}^4 \hat{\alpha}_j(t, \mu, y_0) \right) \right| \\ &\leq \left| y - y_0 - (D^2k(y_0))^{-1} \left(\nabla k(y) + O_{A_0}(\mu \rho^2 + \mu^2 \rho + \mu^{3+\frac{1}{4}}) \right) \right| \\ &\leq O_{A_0}(\rho^2 + \mu^2 \rho + \mu^{3+\frac{1}{4}}). \end{aligned}$$

Hence, we may choose for small μ a radius $0 < \rho = O_{A_0}(\mu^{3+\frac{1}{4}})$ such that F maps U_ρ into itself and $U_\rho \subset B_{\delta_1}(y_0)$. Consequently, the unique fixed-point $\beta(t, \mu)$ must lie in this ball. This ends the proof. $\square$

Lemma 6.2. *Under the assumptions of Lemma 6.1, if moreover $k \in C^6(\mathbb{R}^3)$ and $\|D^6 k\|_\infty \leq A_0$, then we have*

$$\begin{aligned} (\alpha(t, \mu, \beta(t, \mu)))_0 &= -t\mu^3(1 + tk(y_0))^{-\frac{5}{4}} \frac{3^{\frac{3}{4}} 4}{\pi\sqrt{5}} a_0(y_0) \\ &\quad + t\mu^4(1 + tk(y_0))^{-\frac{9}{4}} \frac{\pi 3^{\frac{3}{4}} \sqrt{5}}{30} (a_1(y_0) + ta_2(y_0)) \\ &\quad + t\mu^5(1 + tk(y_0))^{-\frac{5}{4}} \frac{\pi 3^{\frac{3}{4}} \sqrt{5}}{30} a_3(y_0) \\ &\quad + O_{A_0}(t\mu^{5+\frac{1}{2}} + t^2\mu^{4+\frac{1}{4}}), \end{aligned}$$

where $a_i(y_0) = a_i(\theta)$ given in (1.7) and (1.8) with $k_\theta = k(\cdot + y_0)$. If the assumption $k \in C^6(\mathbb{R}^3)$ is dropped then the terms of order higher than 4 in μ have to be replaced by $O_{A_0}(t\mu^{4+\frac{1}{4}})$.

Proof. In view of Lemma 6.1 and because $\nabla k(y_0) = 0$ we may estimate functions of $y := \beta(t, \mu)$ and of $k(y) = k(\beta(t, \mu))$ as follows

$$\begin{aligned} F(y) &= F(y_0) + F'(y_0)(D^2 k(y_0))^{-1} \left(\sum_{j=3}^4 \hat{\alpha}_j(t, \mu, y_0) \right) + O_{A_0}(\mu^{3+\frac{1}{4}}), \\ F(k(y)) &= F(k(y_0)) + O_{A_0}(\mu^4). \end{aligned} \quad (6.1)$$

To prove the claim of the lemma we expand $\alpha(t, \mu, \beta(t, \mu))_0$ according to Lemma 4.8 and use (6.1). For instance we have

$$\begin{aligned} (\vec{\alpha}_2(t, \mu, y))_0 &= -t\mu^2(1 + tk(y_0))^{-\frac{5}{4}} \frac{\pi}{3^{\frac{1}{4}} \sqrt{5}} \Delta k(y) + O_{A_0}(t\mu^6) \\ &= t\mu^4(1 + tk(y_0))^{-\frac{5}{4}} \left(\frac{\pi}{3^{\frac{1}{4}} 2 \sqrt{5}} \nabla \Delta k(y_0) (D^2 k(y_0))^{-1} \nabla (\Delta k(y_0)) \right. \\ &\quad \left. + \mu \frac{3^{\frac{3}{4}} 8}{\pi \sqrt{5}} \nabla \Delta k(y_0) \cdot (D^2 k(y_0))^{-1} \oint \frac{(k(x + y_0) - T_{k(\cdot + y_0), 0}^3(x)) x_i}{|x|^8} \right) \\ &\quad + O_{A_0}(t\mu^{5+\frac{1}{4}}). \end{aligned}$$

If we continue expanding the remaining terms given in Lemma 4.8 the claim follows. $\square$

Lemma 6.3. *Under the assumptions of Lemma 6.1 let $\Delta k(y_0) = 0 = a_0(y_0)$ and define*

$$\gamma(t, \mu) := \frac{1}{t\mu^4} (1 + tk(y_0))^{\frac{9}{4}} \frac{30}{\pi 3^{\frac{3}{4}} \sqrt{5}} (\alpha(t, \mu, \beta(t, \mu)))_0.$$

Then

$$\frac{\partial \gamma(t, \mu)}{\partial t} = a_2(y_0) + O_{A_0}(\mu^{\frac{1}{4}}).$$

Proof. We have

$$\frac{d(\vec{\alpha}(t, \mu, \beta(t, \mu)))_0}{dt} = \frac{\partial(\vec{\alpha})_0}{\partial t} \Big|_{(t, \mu, \beta(t, \mu))} + \frac{\partial(\vec{\alpha})_0}{\partial y} \Big|_{(t, \mu, \beta(t, \mu))} \frac{\partial \beta}{\partial t} \Big|_{(t, \mu)}.$$

The derivatives of $(\vec{\alpha})_0$ are computed in (5.2) and (5.17). In order to compute the derivative of β we use the fact that

$$\tilde{\alpha}(t, \mu, \beta) := ((\vec{\alpha}(t, \mu, \beta))_1, \dots, (\vec{\alpha}(t, \mu, \beta))_3)^T = \vec{0}.$$

By (6.1) and Lemmas 5.1 and 5.3 we have

$$\begin{aligned} \frac{\partial \beta}{\partial t} \Big|_{(t, \mu)} &= - \left(\frac{\partial \tilde{\alpha}}{\partial y} \Big|_{(t, \mu, \beta(t, \mu))} \right)^{-1} \frac{\partial \tilde{\alpha}}{\partial t} \Big|_{(t, \mu, \beta(t, \mu))} \\ &= t^{-1} \mu^{-1} \frac{3^{\frac{1}{4}} \sqrt{5}}{\pi} (1 + tk(y_0))^{\frac{5}{4}} \left((D^2 k(y_0))^{-1} + O_{A_0}(\mu^{\frac{1}{4}}) \right) \\ &\quad \left[\frac{1}{t} (\vec{\alpha}(t, \mu, \beta))_i - O_{A_0} \left(\sum_{j=1}^3 (\vec{\alpha}_j(t, \mu, \beta))_i \right) + t O_{A_0}(\mu^{3+\frac{1}{2}}) \right]_{i=1 \dots 3} \\ &= O_{A_0}(\mu^{2+\frac{1}{2}}), \end{aligned}$$

where we used the fact that as $\vec{\alpha}(t, \mu, \beta)_i \equiv 0$ for $1 \leq i \leq 3$

$$\sum_{j=1}^3 (\vec{\alpha}_j(t, \mu, \beta))_i = O_{A_0}(t\mu^{3+\frac{1}{2}}).$$

From (5.2) we get

$$\frac{\partial(\vec{\alpha})_0}{\partial y} \Big|_{(t, \mu, \beta(t, \mu))} \frac{\partial \beta}{\partial t} \Big|_{(t, \mu)} = O_{A_0}(t\mu^{4+\frac{1}{2}}).$$

Furthermore, by Lemmas 5.3

$$\begin{aligned} \frac{d\vec{\alpha}(t, \mu, \beta)_0}{dt} &= \frac{1}{t} \vec{\alpha}(t, \mu, \beta)_0 + \sum_{j=2}^4 \frac{\partial \vec{\alpha}_j(t, \mu, \beta)_0}{\partial t} - \frac{1}{t} \vec{\alpha}_j(t, \mu, \beta)_0 \\ &\quad + O_{A_0}(t\mu^{4+\frac{1}{4}}). \end{aligned}$$

The definition of γ , (6.1), and Lemma 6.2 yield the claim. $\square$

Lemma 6.4. *Under the assumptions of Lemma 6.3 suppose $a_2(y_0) \neq 0$ and either $a_1(y_0) \neq 0$ or $(a_1(y_0) = 0 \text{ and } a_3(y_0) \neq 0)$. Moreover let*

$$A_0 \geq \begin{cases} |a_1(y_0)|^{-1} + 2|a_1(y_0)||a_2(y_0)|^{-1} + |a_2(y_0)|^{-1} & \text{if } a_1(y_0) \neq 0, \\ |a_2(y_0)|^{-1} + |a_3(y_0)|^{-1} + |a_3(y_0)| & \text{if } a_1(y_0) = 0, \end{cases}$$

and $-\frac{a_1(y_0)}{a_2(y_0)} \in (-B_0, B_1)$. Then there exist $\delta_2 = \delta_2(A_0) > 0$ and a C^1 -function $\tilde{t}$,

$$\tilde{t} : \{\mu : 0 < \mu < \delta_2\} \rightarrow (-B_0, B_1) \setminus \{0\},$$

such that $(\alpha(\tilde{t}(\mu), \mu, \beta(\tilde{t}(\mu), \mu)))_0 \equiv 0$ for all $0 < \mu < \delta_2$ and

$$\tilde{t}(\mu) = -\frac{1}{a_2(y_0)} \begin{cases} a_1(y_0) + O_{A_0}(\mu^{\frac{1}{4}}) & \text{if } a_1(y_0) \neq 0 \\ a_3(y_0)\mu + O_{A_0}(\mu^{1+\frac{1}{4}}) & \text{if } a_1(y_0) = 0. \end{cases} \quad (6.2)$$

Moreover $\tilde{t}$ is unique in the sense that, if $t \in (-B_0, B_1)$ and $0 < \mu < \delta_2$ satisfy $(\alpha(t, \mu, \beta(t, \mu)))_0 = 0$ then $t = \tilde{t}(\mu)$.

Proof. We only sketch the proof, which is similar to the proof of Lemma 6.1. We will apply Banach's fixed-point theorem to the function

$$F_\mu(t) = F(t, \mu) := t - a_2(y_0)^{-1} \gamma(t, \mu),$$

where γ is given in Lemma 6.3. To this end we show that for small μ the map F_μ is a contraction in some ball centered at $-\frac{a_1(y_0)}{a_2(y_0)}$ if $a_1(y_0) \neq 0$ and in $B_r(-\frac{a_3(y_0)}{a_2(y_0)}\mu)$, if $a_1(y_0) = 0$, where

$$0 < r \leq r_0 = r_0(\mu) := \frac{1}{2} \mu \frac{|a_3(y_0)|}{|a_2(y_0)|}.$$

To prove that F_μ is a contraction we may proceed as in Lemma 6.1. We only need the derivative of γ , which is given in Lemma 6.3. $\square$

7. A PRIORI ESTIMATES

We combine the results of Sections 3-6 to prove the C^2 -a priori estimates announced in the introduction.

Theorem 7.1. *Suppose there is $A_0 > 2$ such that $k \in C^5(S^3)$ satisfies*

$D^2 k_\theta(0)$ is invertible, if $\theta \in \mathcal{A} := \{\theta \in S^3 : \nabla k(\theta) = 0 \text{ and } \Delta k(\theta) = 0\}$,

$$(A_0)^{-1} \leq 1 + (1 + A_0^{-1})k(\theta) \leq A_0,$$

$\|k\|_{C^5(S^3)} \leq A_0$, and

$$A_0 \geq \sup\{\|(D^2 k_\theta(0))^{-1}\| : \theta \in \mathcal{A}\}.$$

Thus, $\mathcal{A}$ is discrete and there is $r = r(A_0) > 0$ such that

$$\nabla k(\theta) \neq 0 \text{ for all } \theta \in \bigcup_{\theta_0 \in \mathcal{A}} \overline{B_r(\theta_0)} \setminus \{\theta_0\}.$$

Additionally, assume there is $A_1 > 0$ such that

$$A_1 \geq \sup\{|\Delta k_\theta(0)|^{-1} : |\nabla k(\theta)| \leq A_1^{-1} \text{ and } \theta \in S^3 \setminus \bigcup_{\theta_0 \in \mathcal{A}} B_r(\theta_0)\},$$

$$A_1 \geq \sup\{|a_0(\theta)|^{-1} : \theta \in \mathcal{A} \text{ and } a_0(\theta) \neq 0\}.$$

Denote by M the finite set

$$M := \{\theta \in S^3 : \theta \in \mathcal{A}, a_0(\theta) = 0, \text{ and } a_2(\theta) \neq 0\}.$$

Then for every $\delta > 0$ exists $C = C(\mathcal{A}, A_0, A_1, \delta)$ such that for all

$$t \in (0, 1] \setminus \bigcup_{\theta \in M} B_\delta(-a_1(\theta)/a_2(\theta))$$

and solutions φ_t of (1.5) we have

$$C^{-1} \leq \varphi_t(x) \leq C \text{ and } \|\varphi_t(x)\|_{C^{2,\alpha}(S^3)} \leq C.$$

Proof. Set $I_{\delta,k} := (0, 1] \setminus \bigcup_{y \in M} B_\delta(-a_1(y)/a_2(y))$. To obtain a contradiction, we assume that there are sequences $(k_i) \in C^5(S^3)$, satisfying the assumptions of the theorem with $(\mathcal{A}, A_0, A_1, \delta)$ fixed, and $(t_i, \varphi_{t_i}) \in I_{\delta,k_i} \times C^2(S^3)$ of solutions to (1.5) with $k = k_i$ such that $\|\varphi_{t_i}\|_\infty \rightarrow \infty$ as $i \rightarrow \infty$. Passing to a subsequence we may assume $t_i \rightarrow t_0$ as $i \rightarrow \infty$. By Corollary 3.2 there

are $\theta \in S^3$, $\mu_i \rightarrow 0$ and $y_i \rightarrow 0$ such that u_{t_i} defined by (1.3) in stereographic coordinates $\mathcal{S}_\theta(\cdot)$ solves (1.6) and satisfies

$$\begin{aligned} \tilde{w}_{t_i} &:= u_{t_i} - (1 + t_i(k_i)_\theta(y_i))^{-\frac{1}{4}} z_{\mu_i, y_i} \text{ is orthogonal to } T_{z_{\mu_i, y_i}} Z, \\ \|\tilde{w}_{t_i}\|_{\mathcal{D}^{1,2}(\mathbb{R}^3)} &= o(1). \end{aligned}$$

Using the notation of Lemma 4.2 we have with $k = k_i$

$$0 = f'_{t_i}(u_{t_i}) = f'_{t_i}(z_{\mu_i, y_i} + w_0(t_i, \mu_i, y_i) + \tilde{w}_{t_i}).$$

Consequently, for large i , due to the uniqueness of $\vec{\alpha}$ and w in Lemma 4.2,

$$u_{t_i} = z_{\mu_i, y_i} + w(t_i, \mu_i, y_i, k_i) \text{ and } \vec{\alpha}(t_i, \mu_i, y_i, k_i) = 0,$$

where we added the additional parameter k_i to express the dependence of $\vec{\alpha}$ and w on k_i . From the expansion of $\vec{\alpha}$ in (4.4) we see

$$\lim_{i \rightarrow \infty} \nabla(k_i)_\theta(y_i) = 0 \text{ and } \lim_{i \rightarrow \infty} \Delta(k_i)_\theta(y_i) = 0.$$

As $(\mathcal{A}, A_0, A_1, \delta)$ is fixed, the point θ is in $\mathcal{A}$, hence θ is a nondegenerated critical point of each k_i . We may apply Lemma 6.1 with $k = k_i$ and get for large i

$$y_i = \beta(t_i, \mu_i, k_i),$$

where again the additional parameter k_i denotes the dependence on k_i . From Lemma 6.2 we now get

$$\begin{aligned} 0 &= \frac{1}{t_i \mu_i^3} \left(\vec{\alpha}(t_i, \mu_i, \beta(t_i, \mu_i, k_i), k_i) \right) \\ &= -(1 + t_i k_i(\theta))^{-\frac{5}{4}} \frac{3^{\frac{3}{4}} 4}{\pi \sqrt{5}} a_0(\theta, k_i) \\ &\quad + \mu_i (1 + t_i k_i(\theta))^{-\frac{9}{4}} \frac{\pi 3^{\frac{3}{4}} \sqrt{5}}{30} \left(a_1(\theta, k_i) + t_i a_2(\theta, k_i) \right) \\ &\quad + O(\mu_i^{1+\frac{1}{4}}). \end{aligned} \tag{7.1}$$

Consequently, as $(\mathcal{A}, A_0, A_1, \delta)$ is fixed, $a_0(\theta, k_i) = 0$ for large i .

We observe that $|a_2(\theta, k_i)| \geq A_0^{-4}$ for all i large enough, if not then we get, up to a subsequence,

$$\begin{aligned} |a_1(\theta, k_i)| &\geq |k_i(\theta)|^{-1} \left(\frac{15}{8\pi} \left| \int_{\partial B_1(0)} |D^2(k_i)_\theta(0)(x)^2|^2 \right| - A_0^{-4} \right) \\ &\geq \text{const} A_0^{-3} (1 - A_0^{-2}), \end{aligned}$$

which yields a positive lower bound on $|a_1(\theta, k_i) + t_i a_2(\theta, k_i)|$ contradicting the expansion in (7.1) for i large. Hence from (7.1) we infer

$$\left| t_i + \frac{a_1(\theta, k_i)}{a_2(\theta, k_i)} \right| \leq |a_2(\theta, k_i)|^{-1} O(\mu_i^{\frac{1}{4}}) \leq O(\mu_i^{\frac{1}{4}}),$$

which is impossible for $\delta > 0$. This shows that all solutions φ_t of (1.5) with $t \in I_\delta$ are uniformly bounded. From Harnack's inequality and standard elliptic estimates they are uniformly bounded below by a positive constant and uniformly bounded in $C^{2,\alpha}(S^3)$, which ends the proof. $\square$

From the proof of Theorem 7.1 it is clear that k need only to be in $C^4(S^3)$, but then the constant C will also depend on the modulus of continuity of D^4k .

Proof of Theorems 1.2 and 1.4. If $\theta \in M_+^* \cup M_0^*$ we may apply Lemmas 6.1 and 6.4 with $k = k_\theta$ and $y_0 = 0$. If we set $y(\mu) := \beta(\tilde{t}(\mu), \mu)$ then we have $\tilde{\alpha}(\tilde{t}(\mu), \mu, y(\mu)) = 0$ for all $0 < \mu < \min(\delta_1, \delta_2)$ and $y(\mu) = O(\mu^2)$. From Lemma 4.2 we get that

$$\psi(\mu) := z_{\mu, y(\mu)} + w(\tilde{t}(\mu), \mu, y(\mu))$$

is a solution of (1.6) with $t = \tilde{t}(\mu)$. As $\nabla k_\theta(0) = 0$ and $y(\mu) = O(\mu^2)$ we may use (6.1) to obtain in $\mathcal{D}^{1,2}(\mathbb{R}^3)$

$$\psi(\mu) = (1 + \tilde{t}(\mu)k_\theta(0))^{-\frac{1}{4}}z_{\mu,0} + O(\mu^2).$$

To show that $\psi(\mu)$ is positive for small μ , we note that from Sobolev's inequality $\psi(\mu)^- \rightarrow 0$ in L^6 as $\mu \rightarrow 0$, where $\psi(\mu)^- := \min(\psi(\mu), 0)$. Testing $f_t'(\psi(\mu))$ with $\psi(\mu)^-$ and using Sobolev's inequality we get for some $c(k) > 0$

$$\int |\nabla \psi(\mu)^-|^2 = \int (1 + \tilde{t}(\mu)k_\theta(x))(\psi(\mu)^-)^6 \leq c(k) \left(\int |\nabla \psi(\mu)^-|^2 \right)^3.$$

If $\psi(\mu)^- \neq 0$ for small μ we obtain the contradiction

$$c(k)^{-\frac{1}{2}} \leq \int |\nabla \psi(\mu)^-|^2 = \int (1 + \tilde{t}(\mu)k_\theta(x))(\psi(\mu)^-)^6 \xrightarrow{\mu \rightarrow 0} 0.$$

The C^0 -estimate then follows from elliptic regularity (see [8]). Setting

$$\varphi^\theta(\mu, \cdot) := (\mathcal{R}_\theta)^{-1}(\psi(\mu)) \text{ and } t^\theta(\mu) = \tilde{t}(\mu)$$

yields the existence of the desired curve of solutions.

To prove uniqueness of the curves suppose (t_i, φ_i) blows up at $\theta \in S^3$. If $t_i \in (\delta, 1 + \delta)$ then, as in the proof of Theorem 7.1, we get $\theta \in M_+^*$. Under the assumptions of Theorem 1.4 we already know that $\theta \in M_0^*$. If all but finitely many (t_i, φ_i) lie on the curve corresponding to $\theta \in M_+^* \cup M_0^*$, we are done. Hence we may assume, going to a subsequence if necessary, that none of the (t_i, φ_i) lie on the curve. This is impossible since by Corollary 3.2 and Lemma 4.2 there are μ_i, y_i converging to zero such that for i large

$$\mathcal{R}_\theta(\varphi_i) = z_{\mu_i, y_i} + w(t_i, \mu_i, y_i) \text{ and } \tilde{\alpha}(t_i, \mu_i, y_i) = 0,$$

and thus applying the uniqueness part in Lemmas 6.1 and 6.4 we see that $y_i = \beta(t_i, \mu_i)$ and $t_i = \tilde{t}(\mu_i)$ and the points (t_i, φ_i) have to lie on the curve. $\square$

APPENDIX A. FORMULAS AND INTEGRALS

The *Jacobi* polynomial $\mathcal{P}_j^{(\sigma, \sigma)}$ is defined by

$$\mathcal{P}_j^{(\sigma, \sigma)}(x) := \frac{(-1)^j}{2^j j!} (1 - x^2)^{-\sigma} \frac{d^j}{dx^j} ((1 - x^2)^{\sigma+j}) \quad (\text{A.1})$$

To compute integrals containing Jacobi-polynomials we will use

$$\int_{-1}^1 (1 - \xi^2)^\sigma \mathcal{P}_j^{(\sigma, \sigma)}(\xi) \mathcal{P}_i^{(\sigma, \sigma)}(\xi) d\xi = \frac{2^{2\sigma+1} \Gamma(j + \sigma + 1)^2}{(2j + 2\sigma + 1) j! \Gamma(j + 2\sigma + 1)} \delta_{i,j}, \quad (\text{A.2})$$

and the recurrence relation for $j \in \mathbb{N}_0$

$$\begin{aligned} \xi \mathcal{P}_j^{(\sigma, \sigma)}(\xi) &= \frac{(j+1)(j+2\sigma+1)}{(2j+2\sigma+1)(j+\sigma+1)} \mathcal{P}_{j+1}^{(\sigma, \sigma)}(\xi) + \frac{j+\sigma}{2j+2\sigma+1} \mathcal{P}_{j-1}^{(\sigma, \sigma)}(\xi), \\ \mathcal{P}_{-1}^{(\sigma, \sigma)}(\xi) &:= 0, \quad \mathcal{P}_0^{(\sigma, \sigma)}(\xi) := 1. \end{aligned} \quad (\text{A.3})$$

For a detailed account on *Jacobi* polynomial we refer to [22]. To evaluate integrals of the form

$$\int_0^\infty r^a (1+r^2)^{-b} \mathcal{P}_j^{(\sigma, \sigma)}\left(1 - \frac{2}{1+r^2}\right) dr,$$

we use the following change of coordinates

$$\xi = \frac{r - r^{-1}}{r + r^{-1}} = 1 - \frac{2}{1+r^2}, \quad (\text{A.4})$$

which gives

$$r = \left(\frac{1+\xi}{1-\xi}\right)^{1/2} \text{ and } dr = \left(\frac{1+\xi}{1-\xi}\right)^{-1/2} (1-\xi)^{-2} d\xi,$$

and leads to

$$2^{-b} \int_{-1}^1 (1+\xi)^{\frac{a-1}{2}} (1-\xi)^{b-2-\frac{a-1}{2}} \mathcal{P}_j^{(\sigma, \sigma)}(\xi) d\xi.$$

Moreover, we note that for any $a, b > -1$

$$\int_{-1}^1 (1+\xi)^a (1-\xi)^b d\xi = 2^{a+b+1} \frac{\Gamma(a+1)\Gamma(b+1)}{\Gamma(a+b+2)}. \quad (\text{A.5})$$

This gives for $a > -1$ and $2b - a > 1$

$$\int_0^\infty r^a (1+r^2)^{-b} = \frac{\Gamma(1+(a-1)/2)\Gamma(b-1-(a-1)/2)}{2\Gamma(b)}. \quad (\text{A.6})$$

To compute integrals over $\mathbb{R}^N$ we use polar coordinates. To compute the resulting integrals over $\partial B_1(0)$ we use the following elementary results:

For $\vec{\beta} \in \mathbb{N}_0^N$ we have

$$\int_{\partial B_1(0)} \prod_{i=1}^N x_i^{2\beta_i} = \frac{2 \prod_{i=1}^N \Gamma(\beta_i + \frac{1}{2})}{\Gamma(\frac{N}{2} + \sum_{i=1}^N \beta_i)}. \quad (\text{A.7})$$

Let $m \geq 2$ and P_m be a homogeneous polynomial of order m in $x \in \mathbb{R}^N$. Then

$$\int_{\partial B_1(0)} P_m(x) = \frac{1}{2N + (N+m)(m-2)} \int_{\partial B_1(0)} (\Delta P_m)(x), \quad (\text{A.8})$$

$$\int_{\partial B_1(0)} P_m(x) x_i = \frac{1}{2(N+2) + (N+m+1)(m-3)} \int_{\partial B_1(0)} (\Delta P_m)(x) x_i. \quad (\text{A.9})$$

As $\frac{\partial}{\partial x_i} D_y^\ell k(y)(x)^\ell = \ell \frac{\partial}{\partial y_i} D_y^{\ell-1} k(y)(x)^{\ell-1}$ we see

$$\int_{\partial B_1(0)} \frac{D^{2\ell} k(y)(x)^{2\ell}}{(2\ell)!} = \frac{2\pi^{N/2} \Delta^\ell k(y)}{\Gamma(N/2) \prod_{m=1}^\ell 2m(N+2m-2)}, \quad (\text{A.10})$$

$$\int_{\partial B_1(0)} \frac{D^{2\ell+1} k(y)(x)^{2\ell+1} x_i}{(2\ell+1)!} = \frac{2\pi^{N/2} \frac{\partial}{\partial y_i} \Delta^\ell k(y)}{N\Gamma(N/2) \prod_{m=1}^\ell 2m(N+2m)}. \quad (\text{A.11})$$

Lemma A.1. *Suppose $j \in \mathbb{N}_0$, then*

$$\begin{aligned} \int_0^\infty r^6(1+r^2)^{-5} \left(1 - \frac{2}{1+r^2}\right) \mathcal{P}_j^{(\frac{5}{2}, \frac{5}{2})} \left(1 - \frac{2}{1+r^2}\right) \\ = \frac{\Gamma(\frac{3}{2})\Gamma(\frac{7}{2}+j)(j^2+6j+2)}{2\Gamma(6+j)}. \end{aligned} \quad (\text{A.12})$$

Proof. We use the change of coordinates in (A.4) and obtain

$$\begin{aligned} \int_0^\infty r^4(1+r^2)^{-5} \left(1 - \frac{2}{1+r^2}\right) \mathcal{P}_j^{(\frac{5}{2}, \frac{5}{2})} \left(1 - \frac{2}{1+r^2}\right) \\ = 2^{-5} \int_{-1}^1 (1-\xi^2)^{\frac{5}{2}} \frac{\xi}{(1-\xi)^2} \mathcal{P}_j^{(\frac{5}{2}, \frac{5}{2})}(\xi) \\ = \frac{(-1)^j}{2^{j+5}j!} \int_{-1}^1 \left(\frac{1}{(1-\xi)^2} - \frac{1}{1-\xi} \right) \frac{d^j}{d\xi^j} ((1-\xi^2)^{\frac{5}{2}+j}) \\ = \frac{(-1)^j}{2^{j+5}j!} \int_{-1}^1 \left(\frac{(-1)^j(j+1)!}{(1-\xi)^{2+j}} - \frac{(-1)^jj!}{(1-\xi)^{1+j}} \right) (1-\xi^2)^{\frac{5}{2}+j}. \end{aligned}$$

Now, the claim follows from (A.5). $\square$

Lemma A.2. *Let $\beta_j := \frac{(3+j)j!}{\Gamma(j+\frac{5}{2})}$ for $j \in \mathbb{N}$. Then*

$$\int_0^\infty r^6(1+r^2)^{-7} \left(1 - \frac{2}{1+r^2}\right) \left(\sum_{j=0}^\infty \beta_j \mathcal{P}_j^{(\frac{5}{2}, \frac{5}{2})} \left(1 - \frac{2}{1+r^2}\right) \right)^2 = \frac{1}{288}. \quad (\text{A.13})$$

Proof. We use the change of variable given in (A.4) and the recurrence formula (A.3), applied to $\xi \mathcal{P}_j^{(\frac{5}{2}, \frac{5}{2})}(\xi)$, and get

$$\begin{aligned} \int_0^\infty r^6(1+r^2)^{-7} \left(1 - \frac{2}{1+r^2}\right) \left(\sum_{j=0}^\infty \beta_j \mathcal{P}_j^{(\frac{5}{2}, \frac{5}{2})} \left(1 - \frac{2}{1+r^2}\right) \right)^2 \\ = 2^{-7} \int_{-1}^1 (1-\xi^2)^{\frac{5}{2}} \xi \left(\sum_{j=0}^\infty \beta_j \mathcal{P}_j^{(\frac{5}{2}, \frac{5}{2})}(\xi) \right)^2 \\ = 2^{-7} \sum_{d=0}^\infty \sum_{j=0}^d \beta_j \beta_{d-j} \int_{-1}^1 (1-\xi^2)^{\frac{5}{2}} \xi \mathcal{P}_j^{(\frac{5}{2}, \frac{5}{2})}(\xi) \mathcal{P}_{d-j}^{(\frac{5}{2}, \frac{5}{2})}(\xi) \\ = \sum_{l=0}^\infty \beta_l \beta_{l+1} \frac{\Gamma(l+\frac{7}{2})^2(l+\frac{7}{2})}{\Gamma(l+6)(2l+8)(2l+6)l!} = \frac{1}{4} \sum_{l=0}^\infty \frac{(l+1)!}{(l+5)!} = \frac{1}{288}. \end{aligned}$$

$\square$

REFERENCES

- [1] A. Ambrosetti and M. Badiale. *Homoclinics: Poincaré-Melnikov type results via a variational approach*. Ann. Inst. H. Poincaré Anal. Non Linéaire, **15** (1998), no. 2, 233–252.
- [2] A. Ambrosetti, J. Garcia Azorero and I. Peral. *Perturbation of $\Delta u + u^{(N+2)/(N-2)} = 0$, the scalar curvature problem in $\mathbf{R}^N$, and related topics*. J. Funct. Anal., **165** (1999), no. 1, 117–149.
- [3] T. Aubin. *Some nonlinear problems in Riemannian geometry*. Springer Monographs in Mathematics. Springer-Verlag, Berlin (1998).
- [4] T. Aubin and A. Bahri. *Une hypothèse topologique pour le problème de la courbure scalaire prescrite*. J. Math. Pures Appl. (9), **76** (1997), no. 10, 843–850.
- [5] A. Bahri and J.-M. Coron. *The scalar-curvature problem on the standard three-dimensional sphere*. J. Funct. Anal., **95** (1991), no. 1, 106–172.
- [6] G. Bianchi. *Non-existence and symmetry of solutions to the scalar curvature equation*. Comm. Partial Differential Equations, **21** (1996), no. 1-2, 229–234.
- [7] J.-P. Bourguignon and J.-P. Ezin. *Scalar curvature functions in a conformal class of metrics and conformal transformations*. Trans. Amer. Math. Soc., **301** (1987), no. 2, 723–736.
- [8] H. Brezis and T. Kato. *Remarks on the Schrödinger operator with singular complex potentials*. J. Math. Pures Appl. (9), **58** (1979), no. 2, 137–151.
- [9] L. A. Caffarelli, B. Gidas and J. Spruck. *Asymptotic symmetry and local behavior of semilinear elliptic equations with critical Sobolev growth*. Comm. Pure Appl. Math., **42** (1989), no. 3, 271–297.
- [10] S.-Y. A. Chang, M. J. Gursky and P. C. Yang. *The scalar curvature equation on 2- and 3-spheres*. Calc. Var. Partial Differential Equations, **1** (1993), no. 2, 205–229.
- [11] C.-C. Chen and C.-S. Lin. *Prescribing scalar curvature on S^N . I. A priori estimates*. J. Differential Geom., **57** (2001), no. 1, 67–171.
- [12] W. Chen and C. Li. *Prescribing scalar curvature on S^n* . Pacific J. Math., **199** (2001), no. 1, 61–78.
- [13] V. Felli and M. Schneider. *Perturbation results of critical elliptic equations of Caffarelli-Kohn-Nirenberg type*. Journal of Differential Equations, **191** (2003), 121–142.
- [14] B. Gidas, W. M. Ni and L. Nirenberg. *Symmetry of positive solutions of nonlinear elliptic equations in $\mathbf{R}^n$* . In *Mathematical analysis and applications, Part A*, vol. 7 of *Adv. in Math. Suppl. Stud.*. Academic Press, New York (1981) pages 369–402.
- [15] A. González-López, N. Kamran and P. J. Olver. *Normalizability of one-dimensional quasi-exactly solvable Schrödinger operators*. Comm. Math. Phys., **153** (1993), no. 1, 117–146.
- [16] J. L. Kazdan and F. W. Warner. *Existence and conformal deformation of metrics with prescribed Gaussian and scalar curvatures*. Ann. of Math. (2), **101** (1975), 317–331.

- [17] L. D. Landau and E. M. Lifshitz. *Quantum mechanics: non-relativistic theory. Theoretical Physics, Vol. 3*. Pergamon Press Ltd., London-Paris (1958).
- [18] Y. Y. Li. *Prescribing scalar curvature on S^n and related problems. I*. J. Differential Equations, **120** (1995), no. 2, 319–410.
- [19] Y. Y. Li. *Prescribing scalar curvature on S^n and related problems. II. Existence and compactness*. Comm. Pure Appl. Math., **49** (1996), no. 6, 541–597.
- [20] M. Schneider. *Prescribing scalar curvature on S^3* (2004). Preprint.
- [21] R. Schoen and D. Zhang. *Prescribed scalar curvature on the n -sphere*. Calc. Var. Partial Differential Equations, **4** (1996), no. 1, 1–25.
- [22] G. Szegő. *Orthogonal polynomials*. American Mathematical Society, Providence, R.I. (1975).

RUPRECHT-KARLS-UNIVERSITÄT, MATHEMATISCHES INSTITUT, IM NEUENHEIMER FELD
288, 69120 HEIDELBERG, GERMANY
E-mail address: mschneid@mathi.uni-heidelberg.de